PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6:

C12N 15/40, C07K 14/18, C12Q 1/70, C07K 16/10, G01N 33/576

A2

(11) International Publication Number:

WO 96/13590

(43) International Publication Date:

9 May 1996 (09.05.96)

(21) International Application Number:

PCT/EP95/04155

(22) International Filing Date:

23 October 1995 (23.10.95)

(30) Priority Data:

94870166.9 21 October 1994 (21.10.94) EP
(34) Countries for which the regional or
international application was filed: AT et al.
95870076.7 28 June 1995 (28.06.95) EP

(34) Countries for which the regional or international application was filed:

AT et al.

(71) Applicant (for all designated States except US): INNOGENET-ICS N.V. [BE/BE]; Industricpark Zwijnaarde 7, Box 4, B-9052 Zwijnaarde (BE).

(72) Inventors; and

- (75) Inventors/Applicants (for US only): MAERTENS, Geert [BE/BE]; Zilversparrenstraat 64, B-8310 Brugge (BE). STUYVER, Lieven [BE/BE]; Holestraat 8, B-2400 Mol (BE).
- (74) Common Representative: INNOGENETICS N.V.; Mrs. Ann De Clercq, Industriepark Zwijnaarde 7, Box 4, B-9052 Zwijnaarde (BE).

(81) Designated States: AL, AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, C1, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, LS, MW, SD, SZ, UG).

Published

Without international search report and to be republished upon receipt of that report.

With a request for rectification under Rule 91.1(f).

(54) Title: NEW SEQUENCES OF HEPATITIS C VIRUS GENOTYPES AND THEIR USE AS PROPHYLACTIC, THERAPEUTIC AND DIAGNOSTIC AGENTS

(57) Abstract

The present invention relates to new genomic nucleotide sequences and amino acid sequences corresponding to the coding reglon of these genomes. The invention relates to new HCV types and subtypes sequences which are different from the known HCV types and subtypes sequences. More particularly, the present invention relates to new HCV type 7 sequences, new HCV type 9 sequences, new HCV type 10 and new HCV type 11 sequences. Also, the present invention relates to new HCV type 1 sequences of subtypes 1d, 1e, 1f and 1g; new HCV type 2 sequences of subtypes 2e, 2f, 2g, 2h, 2i, 2k and 2l; new HCV type 3 sequences of subtype 3g, new HCV type 4 sequences of subtypes 4k, 4l and 4m; a process for preparing them, and their use for diagnosis, prophylaxis and therapy. More particularly, the present invention provides new type-specific sequences of the Core, the E1 and the NS5 regions of new HCV types 7, 9, 10 and 11, as well as of new variants (subtypes) of HCV types 1, 2, 3 and 4. These new HCV sequences are useful to diagnose the presence of HCV type 1, and/or type 2, and/or type 3, and/or type 4, and/or type 9, and/or type 10, and/or type 11 genotypes or serotypes in a biological sample. Moreover, the availability of these new type-specific sequences can increase the overall sensitivity of HCV detection and should also prove to be useful for prophylactic and therapeutic purposes.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

					Manritania
AT	Austria	GB	United Kingdom	MR	
AU	Australia	GE	Georgia	MW	Malawi
	Barbados	GN	Guinea	NE	Niger
BB	_	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	17	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgystan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic	SD	Sudan
CF	Central African Republic		of Korea	SE	Sweden
CG	Congo	KR	Republic of Kores	SI	Slovenia
CH	Switzerland	KZ	Kazakhstan	SK	Slovakia
CI	Côte d'Ivoire		Liechtenstein	SN	Senegal
CM	Cameroon	u	Sri Lanka	· TD	Chad
CN	China	ŁK		TG	Togo
CS	Czechoslovakia	LU	Luxembourg	TJ	Tajikistan
CZ	Czech Republic	LV	Larvia	TT	Trinidad and Tobago
DE	Germany	MC	Monaco	UA	Ukraine
DK	Denmark	MD	Republic of Moldova	บร	United States of Americ
ES	Spain	MG	Madagascar	UZ	Uzbekistan
FI	Finland	ML	Muli	. VN	Viet Nam
FR	France	MN	Mongolia	VIN	
GA	Gabon				

10

15

20

25

30

1

NEW SEQUENCES OF HEPATITIS C VIRUS GENOTYPES AND THEIR USE AS PROPHYLACTIC, THERAPEUTIC AND DIAGNOSTIC AGENTS

The invention relates to new sequences of hepatitis C virus (HCV) genotypes and their use as prophylactic, therapeutic and diagnostic agents.

The present invention relates to new genomic nucleotide sequences and amino acid sequences corresponding to the coding region of these genomes. The invention relates to new HCV types and subtypes sequences which are different from the known HCV types and subtypes sequences. More particularly, the present invention relates to new HCV type 7 sequences, new HCV type 9 sequences, new HCV types 10 and new HCV type 11 sequences. Also the present invention relates to new HCV type 1 sequences of subtypes 1d, 1e, 1f and 1g; new HCV type 2 sequences of subtypes 2e, 2f, 2g, 2h, 2i, 2k and 2l; new HCV type 3 sequences of subtype 3g, new HCV type 4 sequences of subtypes 4k, 4l and 4m; a process for preparing them, and their use for diagnosis, prophylaxis and therapy.

The technical problem underlying the present invention is to provide new HCV sequences from untill now unknown HCV types and/or subtypes. More particularly, the present invention provides new type-specific sequences of the Core, the E1 and the NS5 regions of new HCV types 7, 9, 10 and 11, as well as of new variants (subtypes) of HCV types 1, 2, 3 and 4. These new HCV sequences are useful to diagnose the presence of HCV type 1, and/or type 2, and/or type 3, and/or type 4, and/or type 7, and/or type 9, and/or type 10, and/or type 11 genotypes or serotypes in a biological sample. Moreover, the availability of these new type-specific sequences can increase the overall sensitivity of HCV detection and should also prove to be useful for prophylactic and therapeutic purposes.

Hepatitis C viruses (HCV) have been found to be the major cause of non-A, non-B hepatitis. The sequences of cDNA clones covering the complete genome of several prototype isolates have been determined (Kato et al., 1990; Choo et al., 1991; Okamoto et al., 1992). Comparison of these isolates shows that the variability in nucleotide sequences can be used to distinguish at least 2 different genotypes, type 1 (HCV-1 and HCV-J) and type 2 (HC-J6 and HC-J8),

10

15

20

25

30

with an average homology of about 68%. Within each type, at least two subtypes exist (e.g. represented by HCV-1 and HCV-J), having an average homology of about 79%. HCV genomes belonging to the same subtype show average homologies of more than 90% (Okamoto et al., 1992). However, the partial nucleotide sequence of the NS5 region of the HCV-T isolates showed at most 67% homology with the previously published sequences, indicating the existence of yet another HCV type (Mori et al., 1992). Parts of the 5' untranslated region (UR), core, NS3, and NS5 regions of this type 3 have been published, further establishing the similar evolutionary distances between the 3 major genotypes and their subtypes (Chan et al., 1992). Type 4 was subsequently discovered (Stuyver et al., 1993b; Simmonds et al., 1993a; Bukh et al., 1993; Stuyver et al., 1994a). As well as type 5 (Stuyver et al., 1993b; Simmonds et al., 1993c; Bukh et al., 1993; Stuyver et al., 1994b), and type 6 HCV groups (Bukh et al., 1993; Simmonds et al., 1993c). An overview of the present state of the art regarding HCV genotypes is given in Table 3. The nomenclature system proposed by the inventors of the present application has now been accepted by scientists worldwide (Simmonds et al., 1994).

The aim of the present invention is to provide new HCV nucleotide and amino acid sequences enabling the detection of HCV infection.

Another aim of the present infection is to provide new nucleotide and amino acid HCV sequences enabling the classification of infected biological fluids into different serological groups.

Another aim of the present invention is to provide new nucleotide and amino acid HCV sequences ameliorating the overall HCV detection rate.

Another aim of the present invention is to provide new HCV sequences, useful for the design of HCV prophylactic or therapeutic vaccine compositions.

Another aim of the present invention is to provide a pharmaceutical composition consisting of antibodies raised against the polypeptides encoded by these new HCV sequences, for therapy or diagnosis.

All the aims of the present invention are met by the following embodiments of the present invention.

The present invention relates more particularly to an HCV polynucleic acid, having a nucleotide sequence which is unique to a heretofore unidentified HCV type or subtype which is different from HCV subtypes 1a, 1b, 1c, 2a, 2b, 2c, 2d, 3a, 3b,

10

15

20

25

30

3c, 3d, 3e, 3f, 4a, 4b, 4c, 4d, 4e, 4f, 4g, 4h, 4i, 4j, 5a or 6a, with said HCV subtypes being classified as in Table 3 by comparison of a part of the NS5 gene nucleotide sequence spanning positions 7932 to 8271, with said amino acid numbering being shown in Table 1, and with said polynucleic acid containing at least one nucleotide differing from said known HCV nucleotide sequences, or the complement thereof. The sequence of known HCV isolates may be found in any nucleotide sequence database known in the art (such as for instance the EMBL database).

The present invention thus also relates to a polynucleic acid having a nucleotide sequence which is unique to at least one of HCV subtypes 1d, 1e, 1f, 1g, 2e, 2f, 2g, 2h, 2i, 2k, 2l, 3g, 4k, 4l, 4m, 7a, 7c or 7d, with said HCV subtypes being classified as defined above.

The present invention thus also relates to a polynucleic acid having a nucleotide sequence which is unique to at least one of HCV types 9, 10 or 11, with said HCV types being classified as defined above.

It is to be noted that the nucleotide(s) difference in the polynucleic acids of the invention may involve an amino acid difference in the corresponding amino acid sequences encoded by said polynucleic acids. A composition according to the present invention may contain only polynucleic acid sequences or polynucleic acid sequences mixed with any excipient known in the art of diagnosis, prophylaxis or therapy.

According to a preferred embodiment, the present invention relates to a polynucleic acid encoding an HCV polyprotein comprising in its amino acid sequence at least one of the following amino acid residues:

I15, C38, V44, A49, Q43, P49, Q55, A58, S60 or D60, E68 or V68, H70, A71 or Q71 or N71, D72, H81, H101, D106, S110, L130, I134, E135, L140, S148, T150 or E150, Q153, F155, D157, G160, E165, I169, F181, L186, T190, T192 or I192 or H192, I193, A195, S196, R197 or N197 or K197, Q199 or D199 or H199 or N199, F200 or T200, A208, I213, M216 or S216, N217 or S217 or G217 or K217, T218, I219, A222, Y223, I230, W231 or L231, S232 or H232 or A232, Q233, E235 or L235, F236 or T236, F237, L240 or M240, A242, N244, N249, I250 or K250 or R250, A252 or C252, A254, I255 or V255, D256 or M256, E257, E260 or K260, R261, V268, S272 or R272, I285, G290 or F290, A291, A293 or L293

or W293, T294 or A294, S295 or H295, K296 or E296, Y297 or M297, I299 or Y299, I300, S301, P316, S2646, A2648, G2649, A2650, V2652, Q2653, H2656 or L2656, D2657, F2659, K2663 or Q2663, A2667 or V1667, D2677, L2681, M2686 or Q2686 or E2686, A2692 or K2692, H2697, I2707, L2708 or Y2708, A2709, A2719 or M2719, F2727, T2728 or D2728, E2729, F2730 or Y2730, I2741, I2745, V2746 or E2746 or L2746 or K2746, A2748, S2749 or P2749, R2750, E2751, D2752 or N2752 or S2752 or T2752 or V2752 or I2752 or Q2752, S2753 or D2753 or G2753, D2754, A2755, L2756 or Q2756, R2757,

with said notation being composed of a letter representing the amino acid residue by its one-letter code, and a number representing the amino acid numbering according to Kato et al. (1980), as shown in Table 1,

or a part of said polynucleic acid which is unique to at least one of the HCV subtypes or types as defined in Table 5, and which contains at least one nucleotide differing from known HCV nucleotide sequences, or the complement thereof.

15

20

and 119)

5

10

Each of the above-mentioned residues can be found in Figures 2, 4 or 6 showing the new amino acid sequences of the present invention aligned with known sequences of other types or subtypes of HCV for the Core/E1 region.

According to another preferred embodiment, the present invention relates to a polynucleic acid encoding a HCV polyprotein comprising in its amino acid sequence at least one amino acid sequence chosen from the following list:

ARQSDGRSWAQ or ARRSEGRSWAQ as for subtype 1d (SEQ ID NO 107 and 108)

	ERRPEGRSWAQ as for subtype 1e	(SEQ ID NO 109)
	ARRPEGRSWAQ as for subtype 1f	(SEQ ID NO 110)
25	DRRTTGKSWGR as for subtype 2k	(SEQ ID NO 111)
30	DRRATGRSWGR as for subtype 2e	(SEQ ID NO 112)
	DRRATGKSWGR as for subtype 2f	(SEQ ID NO 113)
	VRQPTGRSWGQ as for type 9	(SEQ ID NO 114)
	VRHQTGRTWAQ as for subtype 7a and 7c	(SEQ ID NO 115)
	VRQNQGRTWAQ as for subtype 7d	(SEQ ID NO 116)
	ARRTEGRSWAQ as for type 10	(SEQ ID NO 117)
	VRRTTGRXXXX or VRRTTGRTWAQ as for type 11	(SEQ ID NO 118

	HEVRNASGVYHV or HEVRNASGVYHL as for subtype and 121)	1d (SEQ ID NO 120)
,	YEVHSTTDGYHV as for subtype 1f	(SEQ ID NO 122)	
	VEVKNTSQAYMA as for subtype 2e	(SEQ ID NO 123)	
5	IQVKNNSHFYMA as for subtype 2f	(SEQ ID NO 124)	
	VQVKNTSTMYMA as for subtype 2g	(SEQ ID NO 125)	
	VQVKNTSHSYMV as for subtype 2h	(SEQ ID NO 126)	
	VQVANRSGSYMV as for subtype 2i	(SEQ ID NO 127)	
	VEIKNTXNTYVL or VEIKNTSNTYVL as for subtype 2k		
10	and 129)		
	INYRNVSGIYYV or INYRNTSGIYHV or INYHNTSGIYHI o	r TNYRNVSGIYHV as	
	for subtype 4k	(SEQ ID NO 130,	
	131, 132 or 133)	·	
	QHYRNVSGIYHV as for subtype 41	(SEQ ID NO 134)	
15	IQVKNASGIYHL as for type 9	(SEQ ID NO 135)	
	AHYTNKSGLYHL as for subtype 7c	(SEQ ID NO 136)	
	LNYANKSGLYHL as for subtype 7d	(SEQ ID NO 137)	
٠.	LEYRNASGLYMV as for type 10	(SEQ ID NO 138)	
	IYEMDGMIMHY or IYEMSGMILHA as for subtype 1d	(SEQ ID NO 139	
20	and 140)		
	VYEAKDIILHT as for subtype 1f	(SEQ ID NO 141)	
	VWQLXDAVLHV as for subtype 2e	(SEQ ID NO 142)	
	VWQLRDAVLHV as for subtype 2f	(SEQ ID NO 143)	
	IWQMQGAVLHV as for subtype 2g	(SEQ ID NO 144)	
25	VWQLKDAVLHV as for subtype 2h	(SEQ ID NO 145)	
	VWQLEEAVLHV as for subtype 2i	(SEQ ID NO 146)	
	TWQLXXAVLHV as for subtype 2k	(SEQ ID NO 147)	
	VYEADHHILHL or VYEADHHILAL or VFEADHHILHL as for		
	ID NO 148, 149 and 150)	•	
30	VYESDHHILHL as for subtype 4I	(SEQ ID NO	
	151)		
•	VFEAETMILHL as for type 9	(SEQ ID NO 152)	45h
	VYEAETLILHL as for subtype 7c	(SEQ ID NO	•••

	153)	4050 ID NO 454)
	VYEANGMILHL as for subtype 7d	(SEQ ID NO 154).
	VYEAGDIILHL as for type 10	(SEQ ID NO 155)
	VREDNHLRCWMAL or VRENNSSRCWMAL as for subtype	
5		156 and 157)
•	IREGNISRCWVPL as for subtype 1f	(SEQ ID NO 158)
·	ENSSGRFHCWIPI as for subtype 2e	(SEQ ID NO 159)
	ERSGNRTFCWTAV as for subtype 2f	(SEQ ID NO 160)
	ELQGNKSRCWIPV as for subtype 2g	(SEQ ID NO 162)
10	ERHQNQSRCWIPV as for subtype 2h	(SEQ ID NO 163)
	EWKDNTSRCWIPV as for subtype 2i	(SEQ ID NO 164)
	EREGNSSRCWIPV as for subtype 2k	(SEQ ID NO 165)
	VREGNQSRCWVAL or VRTGNQSRCWVAL or VRVC	SNQSSCWVAL or
•	VRVGNQSRCWVAL or VKEGNHSRCWVAL as for subtype	e 4k
15	(SEQ ID NO 166,	167, 168 or 169)
*	VKTGNTSRCWVAL as for subtype 4I	(SEQ ID NO 170)
	IKAGNESRCWLPV as for type 9	(SEQ ID NO 171)
	VKEGNQSRCWVQA as for subtype 7c	(SEQ ID NO 172)
	VKXXNLTKCWLSA as for subtype 7d	(SEQ ID NO 173)
20	VRSGNTSRCWIPV as for type 10	(SEQ ID NO 174)
	VKNASVPTAA or VKDANVPTAA as for subtype 1d	(SEQ ID NO 175
	and 176)	
•	ARIANAPIDE as for subtype 1f	(SEQ ID NO 177)
	VSKPGALTKG as for subtype 2e	(SEQ ID NO 178)
25	VSRPGALTRG as for subtype 2f	(SEQ ID NO 179)
	VNQPGALTRG as for subtype 2g	(SEQ ID NO 180)
	VSQPGALTRG as for subtype 2h	(SEQ ID NO 181)
	VSQPGALTKG as for subtype 2i	(SEQ ID NO 182)
	VSRPGALTEG as for subtype 2k	(SEQ ID NO 183)
30	APYIGAPLES or APYTAAPLES as for subtype 4k (SEQ	ID NO 184 and 185)
	APILSAPLMS as for subtype 4I	(SEQ ID NO 186)
	VPNSSVPIHG as for type 9	(SEQ ID NO 187) 🦠
	VPNASTPVTG as for subtype 7c	(SEQ ID NO 188)

25

30

7

	VQNASVSIRG as for subtype 7d	(SEQ ID NO 189)
	VKSPCAATAS as for type 10	(SEQ ID NO 190)
	SPRMHHTTQE or SPRLYHTTQE as for subtype 1d (SEQ ID NO 191 and 192)
	TSRRHWTVQD as for subtype 1f	(SEQ ID NO 193)
5	APKRHYFVQE as for subtype 2e	(SEQ ID NO 194)
	SPQYHTFVQE as for subtype 2f	(SEQ ID NO 195)
	SPQHHNFSQD as for subtype 2g	(SEQ ID NO 196)
	SPQHHIFVQD as for subtype 2h	(SEQ ID NO 197)
	SPEHHHFVQD as for subtype 2k	(SEQ ID NO 198)
10	RPRRHWTTQD or RPRRHWTAQD or QPRRHWTTQD	•
	subtype 4k (SEQ ID NO 199, 2	
	QPRRHWTVQD as for subtype 41	(SEQ ID NO 203)
	RPKYHQVTQD as for type 9	(SEQ ID NO 204)
	RPRMHQVVQE as for subtype 7c	(SEQ ID NO 205)
15	RPRMYEIAQD as for subtype 7d	(SEQ ID NO 206)
	RHRQHWTVQD as for type 10	(SEQ ID NO 207)
•		(024 ID NO 201)

or a part of said polynucleic acid which is unique to at least one of the HCV subtypes or types as defined Table 5, and which contains at least one nucleotide differing from known HCV nucleotide sequences, or the complement thereof.

Using the 5' non-coding LiPA system (Stuyver et al., 1993) and a new core LiPA system including multiple probes for subtypes 1a, 1b, 1c, 2a, 2b or 2c derived from the core region (Stuyver et al., 1995), samples from the Benelux, Cameroon, France and Vietnam were selected because of their aberrant reactivities (isolates CAM1078, FR2, FR1, VN4, VN12, VN13, NE98). Some samples were, together with many other samples, sequenced as a control for typing. Sequencing results, however, indicated the discovery of new subtypes (isolates BNL1, BNL2, BNL3, FR4, BNL4, BNL5, BNL6, BNL7, BNL8, BNL9, BNL10, BNL11 and BNL12). Nucleotide sequences in the core and E1 regions which have not yet been reported before, were analyzed in the frame of the invention. Genomic sequences of subtype 1d, 1e, 1f, 1g 2e, 2f, 2g, 2h, 2i, 2k, 2l, 3g, 4k, 4l, 4m, 7a, 7c, 7d and types 9, 10 and 11 isolates are reported for the first time in the present invention. The NS5B region was also analyzed.

The term "polynucleic acid" refers to a single- stranded or double-stranded

nucleic acid sequence which may contain at least 5 contiguous nucleotides in common with the complete nucleotide sequence (e.g. at least 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50, 60, 75 or more contiguous nucleotides). A polynucleic acid which is up till about 100 nucleotides in length is often also referred to as an oligonucleotide. A polynucleic acid may consist of deoxyribonucleotides or ribonucleotides, nucleotide analogues or modified nucleotides, or may have been adapted for therapeutic purposes. A polynucleic acid may also comprise a double stranded cDNA clone which can be used for cloning purposes, or for *in vivo* therapy, or prophylaxis.

10

5

The oligonucleotides according to the present invention, used as primers or probes may also contain or consist of nucleotide analogous such as phosphorothicates (Matsukura et al., 1987), alkylphosphoriates (Miller et al., 1979) or peptide nucleic acids (Nielsen et al., 1991; Nielsen et al., 1993) or may contain interculating agents (Asseline et al., 1984).

15

As most other variations or modifications introduced into the original DNA sequences of the invention these variations will neccissitate adaptions with respect to the conditions under which the oligonucleotide should be used to obtain the required specificty and sensitivity. However the eventual results will be essentially the same as those obtained with the unmodified oligonucleotides.

20

The introduction of these modifications may be advantageous in order to positivily influence characteristics such as hybridization kinetics, reversibility of the hybrid-formation, biological stability of the oligonucleotide molecules, etc.

The polynucleic acids of the invention may be comprised in a composition of any kind. Said composition may be for diagnostic, therapeutic or prophylactic use.

25

The expression "sequences which are unique to an HCV type or subtype" refers to sequences which are not shared by any other type or subtype of HCV, and can thus be used to uniquely detect that HCV type or subtype. Sequence variability is demonstrated in the present invention between the newly found HCV types and subtypes (see Table 5) and the known HCV types and subtypes (see Table 3), and it is therefore from these regions of sequence variability in particular that type- or subtypes-specific polynucleic acids, oligonucleotides, polypeptides and peptides may be obtained. The term type- or subtypes-specific refers to the fact that a sequence is unique to that HCV type or subtype involved.

30

10

15

20

9

The expression "nucleotides corresponding to" refers to nucleotides which are homologous or complementary to an indicated nucleotide sequence or region within a specific HCV sequence.

The term "coding region" corresponds to the region of the HCV genome that encodes the HCV polyprotein. In fact, it comprises the complete genome with the exception of the 5' untranslated region and 3' untranslated region.

The term "HCV polyprotein" refers to the HCV polyprotein of the HCV-J isolate (Kato et al., 1990). The adenine residue at position 330 (Kato et al., 1990) is the first residue of the ATG codon that initiates the long HCV polyprotein of 3010 amino acids in HCV-J and other type 1b isolates, and of 3011 amino acids in HCV-1 and other type 1a isolates, and of 3033 amino acids in type 2 isolates HC-J6 and HC-J8 (Okamoto et al., 1992).

This adenine is designated as position 1 at the nucleic acid level, and this methionine is designated as position 1 at the amino acid level, in the present invention. As type 1a isolates contain 1 extra amino acid in the NS5A region, coding sequences of type 1a and 1b have identical numbering in the Core, E1, NS3, and NS4 region, but will differ in the NS5B region as indicated in Table 1. Type 2 isolates have 4 extra amino acids in the E2 region, and 17 or 18 extra amino acids in the NS5 region compared to type 1 isolates, and will differ in numbering from type 1 isolates in the NS3/4 region and NS5b regions as indicated in Table 1. Similar insertions compared with type 1 (but of a different size) can also be observed in type 3a sequences which affect the numbering of type 3a amino acids accordingly. Other insertions or deletions may be readily observed in type1, 2, 3, 4, 5, 6, 7, 8, 9, 10 and 11 sequences after alignment withknown HCV sequences.

25

TABLE 1

	Region	Positions described in the present invention*	Positions described for HCV-J (Kato et al., 1990)	Positions described for HCV-1 (Choo et al., 1991)	Positions described for HC-J6, HC- JB (Okamoto et al., 1992)
--	--------	---	---	---	---

Nucleotides	NS5B	8023/8235 7932/8271	8352/8564 8261/8600	8026/8238 7935/8274	8433/8645 8342/8681
		coding region of present invention	330/9359	1/9033	342/9439
Amino Acids	NS5B	2675/2745 2645/2757	2675/2745 2645/2757	2676/2746 2646/2758	2698/2768 2668/2780

Table 1:

5

10

Comparison of the HCV nucleotide and amino acid numbering system used in the present invention (*) with the numbering used for other prototype isolates. For example, 8352/8564 indicates the region designated by the numbering from nucleotide 8352 to nucleotide 8564 as described by Kato et al. (1990). Since the numbering system of the present invention starts at the polyprotein initiation site, the 329 nucleotides of the 5' untranslated region described by Kato et al. (1990) have to be substracted, and the corresponding region is numbered from nucleotide 8023 ('8352-329') to 8235 ('8564-329').

The term "genotype" as used in the present invention refers to both types and/or subtypes.

The term "HCV type" corresponds to a group of HCV isolates of which the 15 complete genome shows more than 73% preferably more than 74% homology at the nucleic acid level, or of which the NS5 region between nucleotide positions 7932 and 8271 shows more than 75.4% homology at the nucleic acid level, or of which the complete HCV polyprotein shows more than 78% homology at the amino acid level, or of which the NS5 region between amino acids at positions 2645 and 2757 20 shows more than 80% homology at the amino acid level, to polyproteins of the other isolates of the group, with said numbering beginning at the first ATG codon or first methionine of the long HCV polyprotein of the HCV-J isolate (Kato et al., 1990). Isolates belonging to different types of HCV exhibit homologies, over the complete genome, of less than 74%, preferably less than 73%, at the nucleic acid level and 25 less than 78% at the amino acid level. Isolates belonging to the same type usually

10

15

20

25

30

show homologies of about 90 to 99% at the nucleic acid level and 95 to 96% at the amino acid level when belonging to the same subtype, and those belonging to the same type but different subtypes preferably show homologies of about 76% to 82% (more particularly of about 77% to 80%) at the nucleic acid level and 85-86% at the amino acid level.

More preferably the definition of HCV types is concluded from the classification of HCV isolates according to their nucleotide distances calculated as detailed below:

- (1) based on phylogenetic analysis of nucleic acid sequences in the NS58 region between nucleotides 7935 and 8274 (Choo et al., 1991) or 8261 and 8600 (Kato et al., 1990) or 8342 and 8681 (Okamoto et al., 1991), isolates belonging to the same HCV type show nucleotide distances of less than 0.34, usually less than 0.33, and more usually of less than 0.32, and isolates belonging to the same subtype show nucleotide distances of less than 0.135, usually of less than 0.13, and more usually of less than 0.125, usually ranging between 0.0003 and 0.1151, and consequently isolates belonging to the same type but different subtypes show nucleotide distances ranging from 0.135 to 0.34, usually ranging from 0.1384 to 0.2977, and more usually ranging from 0.15 to 0.32, and isolates belonging to different HCV types show nucleotide distances greater than 0.34, usually greater that 0.35, and more usually of greater than 0.358, more usually ranging from 0.3581 to 0.6670.
- (2) based on phylogenetic analysis of nucleic acid sequences in the core/E1 region between nucleotides 378 and 957, isolates belonging to the same HCV type show nucleotide distances of less than 0.38, usually of less than 0.37, and more usually of less than 0.364, and isolates belonging to the same subtype show nucleotide distances of less than 0.17, usually of less than 0.16, and more usually of less than 0.15, more usually less than 0.135, more usually less than 0.134, and consequently isolates belonging to the same type but different subtypes show nucleotide distances ranging from 0.15 to 0.38, usually ranging from 0.16 to 0.37, and more usually ranging from 0.17 to 0.36, more usually ranging from 0.133 to 0.379, and isolates belonging to different HCV types show nucleotide distances greater than 0.34, 0.35, 0.36, usually more than 0.365, and more usually of greater than 0.37,

Table 2: Molecular evolutionary distances

Region	Core/E1	E1	NS5B	NS5B
	579 bp	384 bp	340 bp	222 bp
lsolates	0.0017 - 0.1347	0.0026 - 0.2031	0.0003 - 0.1151	0.000 - 0.1323
	(0.0750 ± 0.0245)	(0.0969 ± 0.0289)	(0.0637 ± 0.0229)	(0.0607 ± 0.0205)
Subtypes	0.1330 - 0.3794	0.1645 - 0.4869	0.1384 - 0.2977	0.117 - 0.3538
	(0.2786 ± 0.0363)	. (0.3761 ± 0.0433)	(0.2219 ± 0.0341)	(0.2391 ± 0.0399)
Турез	0.3479 - 0.6306	0.4309 - 0.9561	0.3581 - 0.6670	0.3457 - 0.7471
	(0.4703 ± 0.0525)	(0.6308 ± 0.0928)	(0.4994 ± 0.0495)	(0.5295 ± 0.0627)

Table 2

Figures created by the PHYLIP program DNADIST are expressed as minimum to maximum (average <u>+</u> standard deviation). Phylogenetic distances for isolates belonging to the same subtype ('isolates'), to different subtypes of the same type ('subtypes'), and to different types ('types') are given.

10

5

In a comparative phylogenetic analysis of available sequences, ranges of molecular evolutionary distances for different regions of the genome were calculated, based on 19,781 pairwise comparisons by means of the DNADIST program of the phylogeny inference package PHYLIP version 3.5c (Felsenstein, 1993). The results are shown in Table 2 and indicate that although the majority of distances obtained in each region fit with classification of a certain isolate, only the ranges obtained in the 340bp NS5B-region are non-overlapping and therefore conclusive. However, as was performed in the present invention, it is preferable to obtain sequence information from at least 2 regions before final classification of a given isolate.

20

15

Designation of a number to the different types of HCV and HCV nomenclature is based on chronological discovery of the different types. The numbering system used in the present invention might still fluctuate according to international conventions or guidelines. For example, "type 4" might be changed into "type 5" or "type 6". Also the arbitrarily chosen border distances between types and subtypes and isolates may still be subject to change according to international guidelines or

10

15

20

conventions. Therefore types 7a, 8a, 8b, 9a may for example be designated 6b, 6c, 6d, and 6d in the future; and type 10a which shows relatedness with genotype 3 may be denoted 3g instead of 10a.

The term "subtype" corresponds to a group of HCV isolates of which the complete polyprotein shows a homology of more than 90% both at the nucleic acid and amino acid levels, or of which the NS5 region between nucleotide positions 7932 and 8271 shows a homology of more than 90% at the nucleic acid level to the corresponding parts of the genomes of the other isolates of the same group, with said numbering beginning with the adenine residue of the initiation codon of the HCV polyprotein. Isolates belonging to the same type but different subtypes of HCV show homologies of more than 74% at the nucleic acid level and of more than 78% at the amino acid level.

It is to be understood that extremely variable regions such as the E1, E2 and NS4 regions will exhibit lower homologies than the average homology of the complete genome of the polyprotein.

Using these criteria, HCV isolates can be classified into at least 11 types. Several subtypes can clearly be distinguished in types 1, 2, 3, 4 and 7: 1a, 1b, 1c, 1d, 1e, 1f, 1g, 2a, 2b, 2c, 2d, 2e, 2f, 2g, 2h, 2i, 2k, 2l, 3a, 3b, 3c, 3d, 3f, 3g, 4a, 4b, 4c, 4d, 4e, 4f, 4g, 4h, 4i, 4j, 4k, 4l, 4m, 7a, 7c, and 7d based on homologies of the 5' UR and coding regions. An overview of most of the reported isolates and their proposed classification according to the typing system of the present invention as well as other proposed classifications is presented in Table 3.

Table 3

				HCV	CLASSIFICA	ATION
25		OKA- MOTO	MORI	СНА	NAKAO	PROTOTYPE
	la	1	I	Pt	GI	HCV-1, HCV-H, HC-JI
	1 b	II	II	KI .	GII	HCV-J, HCV-BK, HCV-T, HC-JK1, HC-J4, HCV-CHINA
	Ιc					HC-G9
	2а	III	· III	K2a	GIII	HC-J6
30	2b	IV	IV	К2ь	GIII	HC-J8

	2c					S83, ARG6, ARG8, I10, T983
	2 d					NE92
	3a	V	v	K3	GIV	BR36, BR56, HD10, N2L1, BR33, Ta, E-b1
5	3Ь		VI	К3	GIV	HCV-TR, Tb, NE137
	3c					NE48
	3d					NE274
	3 e					NE145
	3 f					NE125
10	4a					Z4, GB809-4
.0	4b					Z1
	4c					GB116, GB358, GB215, Z6, Z7
	4d					DK13
	4e					GB809-2, CAM600, CAM736
15	4f					CAM622, CAM627
13	4g					GB549
	4h					GB438
	4i				,	CAR4/1205
						CAR1/905
	4j				GV	SA3, SA4, SA1, SA7, SA11, BE95
20	5a				, 0,	HK1, HK2, HK3, HK4, VN11
	6a					FIRT, IIRZ, IIRZ, IIRZ, IIRZ,

Table 3 Overview of the known HCV types and subtypes classified according to the different authors.

The term "complement" refers to a nucleotide sequence which is complementary to an indicated sequence and which is able to hybridize to the indicated sequences.

The composition of the invention can comprise many combinations. By way of example, the composition of the invention can comprise:

- two (or more) nucleic acids from the same region or,
- two nucleic acids (or more), respectively from different regions, for the same isolate or for different isolates,
 - or nucleic acids from the same regions and from at least two different regions

10

15

20

25

30

(for the same isolate or for different isolates).

The present invention relates particularly to a polynucleic acid as defined above having a sequence selected from any of SEQ ID NO 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87, 89, 91, 93, 95, 97, 99, 101, 103 to 105, or a part of said polynucleic acid which is unique to any of the HCV subtypes or types as defined in Table 5, and which contains at least one nucleotide differing from known HCV polynucleic acids, or the complement thereof.

The present invention relates more particularly to a polynucleic acid as defined above, which codes for the 5' UR, the Core/E1, the NS4 or the NS5B region or a part thereof.

More particularly, the present invention relates to a polynucleic acid as defined above which is a cDNA sequence.

Also included within the present invention are sequence variants of the polynucleic acids as selected from any of the nucleotide sequences as given in any of the above given SEQ ID numbers with said sequence variants containing either deletion and/or insertions of one or more nucleotides, especially insertions or deletions of 1 or more codons, mainly at the extremities of oligonucleotides (either 3' or 5'), or substitutions of some non-essential nucleotides (i.e. nucleotides not essential to discriminate between different genotypes of HCV) by others (including modified nucleotides an/or inosine), for example, a type 1 or 2 sequence might be modified into a type 7 sequence by replacing some nucleotides of the type 1 or 2 sequence with type-specific nucleotides of type 7 as shown in for instance Figure 1 and 2.

Particularly preferred variant polynucleic acids of the present invention include also sequences which hybridise under stringent conditions with any of the polynucleic acid sequences of the present invention. Particularly, sequences which show a high degree of homology (similarity) to any of the polynucleic acids of the invention as described above. Particularly sequences which are at least 80%, 85%, 90%, 95% or more homologous to said polynucleic acid sequences of the invention. Preferably said sequences will have less than 20%, 15%, 10%, or 5% variation of the original nucleotides of said polynucleic acid sequence.

Polynucleic acid sequences according to the present invention which are

10

15

20

25

30

homologous to the sequences as represented by a SEQ ID NO can be characterized and isolated according to any of the techniques known in the art, such as amplification by means of sequence-specific primers, hybridization with sequence-specific probes under more or less stringent conditions, serological screening methods or via the LiPA typing system.

Other preferred variant polynucleic acids of the present invention include sequences which are redundant as a result of the degeneracy of the genetic code compared any of the above-given polynucleic acids of the present invention. These variant polynucleic acid sequences will thus encode the same amino acid sequence as the polynucleic acids they are derived from.

Also included within the scope of the present invention are 5' non-coding region sequences which can be readily obtained from type 1 subtype 1d, 1e, 1f or 1g isolates; type 2 subtype 2e, 2f, 2g, 2h, 2i, 2k or 2l isolates; type 3 subtype 3g isolates; type 4 subtype 4k, 4l or 4m isolates; type 7 subtype 7a, 7c or 7d isolates, type 9, type 10 or type 11 isolates discribed herein. Such sequences may contain type or subtype-specific motifs which can be employed for type and/or subtype-specific hybridization assays, e.g. such as described by Stuyver et al. (1993).

Polynucleic acid sequences of the genomes indicated above from regions not yet depicted in the present examples, figures and sequence listing can be obtained by any of the techniques known in the art, such as amplification techniques using suitable primers from the sequences of these new genomes given in Figure 1 of the present invention.

The present invention also relates to an oligonucleotide primer comprising part of a polynucleic acid as defined above, with said primer being able to act as a primer for specifically amplifying the nucleic acid of a certian HCV isolate belonging to the genotype from which the primer is derived.

The term "primer" refers to a single stranded DNA oligonucleotide sequence capable of acting as a point of initiation for synthesis of a primer extension product which is complementary to the nucleic acid strand to be copied. The length and the sequence of the primer must be such that they allow to prime the synthesis of the extension products. Preferably the primer is about 5-50 nucleotides. Specific length and sequence will depend on the complexity of the required DNA or RNA targets, as well as on the conditions of primer use such as temperature and ionic strength.

10

15

20

25

30

The fact that amplification primers do not have to match exactly with corresponding template sequence to warrant proper amplification is amply documented in the literature (Kwok et al., 1990).

The amplification method used can be either polymerase chain reaction (PCR; Saiki et al., 1988), ligase chain reaction (LCR; Landgren et al., 1988; Wu & Wallace, 1989; Barany, 1991), nucleic acid sequence-based amplification (NASBA; Guatelli et al., 1990; Compton, 1991), transcription-based amplification system (TAS; Kwoh et al., 1989), strand displacement amplification (SDA; Duck, 1990; Walker et al., 1992) or amplification by means of Qß replicase (Lizardi et al., 1988; Lomeli et al., 1989) or any other suitable method to amplify nucleic acid molecules using primer extension. During amplification, the amplified products can be conveniently labelled either using labelled primers or by incorporating labelled nucleotides. Labels may be isotopic (³²P, ³⁵S, etc.) or non-isotopic (biotin, digoxigenin, etc.). The amplification reaction is repeated between 20 and 70 times, advantageously between 25 and 45 times.

The present invention also relates to an oligonucleotide probe comprising part of a polynucleic acid as defined above, with said probe being able to act as a hybridization probe for specific detection and/or classification into types and/or subtypes of an HCV nucleic caid containing said nucleotide sequence, with said probe being possibly labelled or attached to a solid substrate.

The term "probe" refers to single stranded sequence-specific oligonucleotides which have a sequence which is complementary to the target sequence of the HCV genotype(s) to be detected.

Preferably, these probes are about 5 to 50 nucleotides long, more preferably from about 10 to 25 nucleotides.

The term "solid support" can refer to any substrate to which an oligonucleotide probe can be coupled, provided that it retains its hybridization characteristics and provided that the background level of hybridization remains low. Usually the solid substrate will be a microtiter plate, a membrane (e.g. nylon or nitrocellulose) or a microsphere (bead). Prior to application to the membrane or fixation it may be convenient to modify the nucleic acid probe in order to facilitate fixation or improve the hybridization efficiency. Such modifications may encompass homopolymer tailing, coupling with different reactive groups such as aliphatic

10

15

groups, $\mathrm{NH_2}$ groups, SH groups, carboxylic groups, or coupling with biotin or haptens.

The present invention also relates to a diagnostic kit for use in determining the genotype of HCV, said kit comprising a primer as defined above.

The present invention also relates to a diagnostic kit for use in determining the genotype of HCV, said kit comprising a probe as defined above.

The present invention also relates to a diagnostic kit as defined above, wherein said probe(s) is(are) attached to a solid substrate.

The present invention also relates to a diagnostic kit as defined above, wherein a range of said probes is attached to specific locations on a solid substrate.

The present invention also relates to a diagnostic kit as defined above, wherein said solid support is a membrane strip and said probes are coupled to the membrane in the form of parallel lines.

The present invention also relates to a method for the detection of HCV nucleic acids present in a biological sample, comprising:

- (i) possibly extracting sample nucleic acid,
- (ii) amplifying the nucleic acid with at least one primer as defined above,
- (iii) detecting the amplified nucleic acids.

The present invention also relates to a method for the detection of HCV nucleic acids present in a biological sample, comprising:

- (i) possibly extracting sample nucleic acid,
- (ii) possibly amplifying the nucleic acid with at least one primer as defiend above, or with a universal HCV primer,
- hybridizing the nucleic acids of the biological sample, possibly under denatured conditions, at appropriate conditions with one or more probes as defined above, with said probes being preferably attached to a solid substrate,
 - (iv) possibly washing at appropriate conditions,
 - (v) detecting the hybrids formed.
- The present invention also relates to a method for detecting the presence of one or more HCV genotypes present in a biological sample, comprising:
 - (i) possibly extracting sample nucleic acid,
 - (ii) specifically amplifying the nucleic acid with at least one primer as defined

10

15

20

25

30

above,

(iii) detecting said amplified nucleic acids.

The present invention also relates to a method for detecting the presence of one or more HCV genotypes present in a biological sample, comprising:

- (i) possibly extracting sample nucleic acid,
- (ii) possibly amplifying the nucleic acid with at least one primer as defined above or with a universal HCV primer,
- hybridizing the nucleic acids of the biological sample, possibly under denatured conditions, at appropriate conditions with one or more probes as defined above, with said probes being preferably attached to a solid substrate,
- (iv) possibly washing at appropriate conditions,
- (v) detecting the hybrids formed,
- (vi) inferring the presence of one or more HCV genotypes present from the observed hybridization pattern.

The present invention also relates to a method as defined above, wherein said probes are further characterized as defined above.

The present invention also relates to a method as defined above, wherein said nucleic acids are labelled during or after amplification.

Preferably, this technique could be performed in the 5' non-coding, Core or NS5B region.

The term "nucleic acid" can also be referred to as analyte strand and corresponds to a single- or double-stranded nucleic acid molecule. This analyte strand is preferentially positive- or negative stranded RNA, cDNA or amplified cDNA.

The term "biological sample" refers to any biological sample (tissue or fluid) containing HCV nucleic acid sequences and refers more particularly to blood serum or plasma samples.

The term "universal HCV primer" refers to oligonucleotide sequences complementary to any of the conserved regions of the HCV genome.

The expression "appropriate" hybridization and washing conditions are to be understood as stringent and are generally known in the art (e.g. Maniatis et al., Molecular Cloning: A Laboratory Manual, New York, Cold Spring Harbor Laboratory, 1982).

10

15

20

25

30

However, according to the hybridization solution (SSC, SSPE, etc.), these probes should be hybridized at their appropriate temperature in order to attain sufficient specificity.

The term "labelled" refers to the use of labelled nucleic acids. This may include the use of labelled nucleotides incorporated during the polymerase step of the amplification such as illustrated by Saiki et al. (1988) or Bej et al. (1990) or labelled primers, or by any other method known to the person skilled in the art.

The process of the invention comprises the steps of contacting any of the probes as defined above, with one of the following elements:

 either a biological sample in which the nucleic acids are made available for hybridization,

or the purified nucleic acids contained in the biological sample

or a single copy derived from the purified nucleic acids,

 or an amplified copy derived from the purified nucleic acids, with said elements or with said probes being attached to a solid substrate.

The expression "inferring the presence of one or more HCV genotypes present from the observed hybridization pattern" refers to the identification of the presence of HCV genomes in the sample by analyzing the pattern of binding of a panel of oligonucleotide probes. Single probes may provide useful information concerning the presence or absence of HCV genomes in a sample. On the other hand, the variation of the HCV genomes is dispersed in nature, so rarely is any one probe able to identify uniquely a specific HCV genome. Rather, the identity of an HCV genotype may be inferred from the pattern of binding of a panel of oligonucleotide probes, which are specific for (different) segments of the different HCV genomes. Depending on the choice of these oligonucleotide probes, each known HCV genotype will correspond to a specific hybridization pattern upon use of a specific combination of probes. Each HCV genotype will also be able to be discriminated from any other HCV genotype amplified with the same primers depending on the choice of the oligonucleotide probes. Comparison of the generated pattern of positively hybridizing probes for a sample containing one or more unknown HCV sequences to a scheme of expected hybridization patterns, allows one to clearly infer the HCV genotypes present in said sample.

The present invention thus relates to a method as defined above, wherein one SUBSTITUTE SHEET (RULE 26)

or more hybridization probes are selected from any of SEQ ID NO 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87, 89, 91, 93, 95, 97, 99, 101, 103 or 105 or sequence variants thereof as defined above.

5

In order to distinguish the amplified HCV genomes from each other, the target polynucleic acids are hybridized to a set of sequence-specific DNA probes targetting HCV genotypic regions (unique regions) located in the HCV polynucleic acids.

Most of these probes target the most type- or subtype-specific regions of HCV genotypes, but some can be caused to hybridize to more than one HCV genotype.

10

According to the hybridization solution (SSC, SSPE, etc.), these probes should be stringently hybridized at their appropriate temperature in order to attain sufficient specificity. However, by slightly modifying the DNA probes, either by adding or deleting one or a few nucleotides at their extremities (either 3' or 5'), or substituting some non-essential nucleotides (i.e. nucleotides not essential to discriminate between types) by others (including modified nucleotides or inosine) these probes or variants thereof can be caused to hybridize specifically at the same hybridization conditions (i.e. the same temperature and the same hybridization solution). Also changing the amount (concentration) of probe used may be beneficial to obtain more specific hybridization results. It should be noted in this context, that probes of the same length, regardless of their GC content, will hybridize specifically at approximately the same temperature in TMACI solutions (Jacobs et al., 1988).

20

15

Suitable assay methods for purposes of the present invention to detect hybrids formed between the oligonucleotide probes and the nucleic acid sequences in a sample may comprise any of the assay formats known in the art, such as the conventional dot-blot format, sandwich hybridization or reverse hybridization. For example, the detection can be accomplished using a dot blot format, the unlabelled amplified sample being bound to a membrane, the membrane being incorporated with at least one labelled probe under suitable hybridization and wash conditions, and the presence of bound probe being monitored.

30

25

An alternative and preferred method is a "reverse" dot-blot format, in which the amplified sequence contains a label. In this format, the unlabelled oligonucleotide probes are bound to a solid support and exposed to the labelled sample under appropriate stringent hybridization and subsequent washing conditions. It is to be

457

10

15

20

25

30

understood that also any other assay method which relies on the formation of a hybrid between the nucleic acids of the sample and the oligonucleotide probes according to the present invention may be used.

According to an advantageous embodiment, the process of detecting one or more HCV genotypes contained in a biological sample comprises the steps of contacting amplified HCV nucleic acid copies derived from the biological sample, with oligonucleotide probes which have been immobilized as parallel lines on a solid support.

According to this advantageous method, the probes are immobilized in a Line Probe Assay (LiPA) format. This is a reverse hybridization format (Saiki et al., 1989) using membrane strips onto which several oligonucleotide probes (including negative or positive control oligonucleotides) can be conveniently applied as parallel lines.

The invention thus also relates to a solid support, preferably a membrane strip, carrying on its surface, one or more probes as defined above, coupled to the support in the form of parallel lines.

The LiPA is a very rapid and user-friendly hybridization test. Results can be read after 4 hours, after the start of the amplification. After amplification during which usually a non-isotopic label is incorporated in the amplified product, and alkaline denaturation, the amplified product is contacted with the probes on the membrane and the hybridization is carried out for about 1 to 1,5 h hybridized polynucleic acid is detected. From the hybridization pattern generated, the HCV type can be deduced either visually, but preferably using dedicated software. The LiPA format is completely compatible with commercially available scanning devices, thus rendering automatic interpretation of the results very reliable. All those advantages make the LiPA format liable for the use of HCV detection in a routine setting. The LiPA format should be particularly advantageous for detecting the presence of different HCV genotypes.

The present invention also relates to a method for detecting and identifying novel HCV genotypes, different from the known HCV genomes, comprising the steps of:

- determining to which HCV genotype the nucleotides present in a biological sample belong, according to the process as defined above,
- in the case of observing a sample which does not generate a hybridization

10

15

20

25

30

pattern compatible with those defined in Table 3, sequencing the portion of the HCV genome sequence corresponding to the aberrantly hybridizing probe of the new HCV genotype to be determined.

The present invention also relates to a method for preparing a polynucleic acid according to the present invention. These methods include any method known in the art for preparing polynucleic acids (e.g. the phosphodiester method for synthesizing oligonucleotides as described by Agarwal et al. 1972, Agnew. Chem. Int. Ed. Engl. 11:451, the phosphotriester method of Hsiung et al. 1979, Nucleic Acid Res. 6:1371, or the automated diethylphosphoramidite method of Baeucage et al. 1981, Tetrahedron Letters 22:1859-1862.). Alternatively, the polynucleic acids of the present invention may be isolated fragments of naturally occuring or cloned DNA or RNA. In addition, the oligonucleotides according to the present invention may be synthesized automatically on commercial instruments sold by a variety of manufacturers.

The present invention particularly also relates to a polypeptide having an amino acid sequence encoded by a polynucleic acid as defined above, or a part thereof which is unique to at least one of the HCV subtypes or types as defined in Table 5, and which contains at least one amino acid differing from any of the known HCV types or subtypes, or an analog thereof being substantially homologous and biologically equivalent.

The term 'polypeptide' refers to a polymer of amino acids and does not refer to a specific length of the product; thus, peptides, oligopeptides, and proteins are included within the definition of polypeptide. This term also does not refer to or exclude post-expression modifications of the polypeptide, for example, glycosylations, acetylations, phosphorylations and the like. Included within the definition are, for example, polypeptides containing one or more analogues of an amino acid (including, for example, unnatural amino acids, PNA, etc.), polypeptides with substituted linkages, as well as other modifications known in the art, both naturally occurring and non-naturally occurring.

The term "unique" is referred above.

By "biologically equivalent" as used throughout the specification and claims, it is meant that the compositions are immunogenically equivalent to the proteins (polypeptides) or peptides of the invention as defined above and below.

10

15

20

25

30

By "substantially homologous" as used throughout the ensuing specification and claims to describe proteins and peptides, it is meant a degree of homology in the amino acid sequence to the proteins or peptides of the invention. Preferably the degree of homology is in excess of 90, preferably in excess of 95, with a particularly preferred group of proteins being in excess of 99 homologous with the proteins or peptides of the invention.

The term "analog" as used throughout the specification or claims to describe the proteins or peptides of the present invention, includes any protein or peptide having an amino acid residue sequence substantially identical to a sequence specifically shown herein in which one or more residues have been conservatively substituted with a biologically equivalent residue. Examples of conservative substitutions include the substitution of one-polar (hydrophobic) residue such as isoleucine, valine, leucine or methionine for another, the substitution of one polar (hydrophillic) residue for another such as between arginine and lysine, between glutamine and asparagine, between glycine and serine, the substitution of one basic residue such as lysine, arginine or histidine for another, or the substitution of one acidic residue, such as aspartic acid or glutamic acid for another. Examples of allowable mutations according to the present inevntion can be found in Table 4.

The phrase "conservative substitution" also includes the use of a chemically derivatized residue in place of a non-derivatized residue provided that the resulting protein or peptide is biologically equivalent to the protein or peptide of the invention.

"Chemical derivative" refers to a protein or peptide having one or more residues chemically derivatized by reaction of a functional side group. Examples of such derivatized molecules, include but are not limited to, those molecules in which free amino groups have been derivatized to form amine hydrochlorides, p-toluene sulfonyl groups, carbobenzoxy groups, t-butyloxycarbonyl groups, chloracetyl groups or formyl groups. Free carboxyl groups may be derivatized to form salts, methyl and ethyl esters or other types of esters or hydrazides. Free hydroxyl groups may be derivatized to form O-acyl or O-alkyl derivatives. The imidazole nitrogen of histidine may be derivatized to form N-imbenzylhistidine. Also included as chemical derivatives are those proteins or peptides which contain one or more naturally-occurring amino acid derivatives of the twenty standard amino acids. For examples: 4-hydroxyproline may be substituted for proline; 5-hydroxylysine may be substituted for lysine; 3-

10

15

20

25

methylhistidine may be substituted for histidine; homoserine may be substituted for senine; and ornithine may be substituted for lysine. The proteins or peptides of the present invention also include any protein or peptide having one or more additions and/or deletions or residues relative to the sequence of a peptide whose sequence is shown herein, so long as the peptide is biologically equivalent to the proteins or peptides of the invention.

It is to be noted that, at the level of the amino acid sequence, at least one amino acids difference (with respect to known HCV amino acid sequences) is sufficient to be part of the invention, which means that the polypeptides of the invention correspond to polynucleic acids having at least one nucleotide difference (with known HCV polynucleic acid sequences) involving an amino acid difference in the encoded polyprotein.

As the NS4 and the Core regions are known to contain several epitopes, for example characterized in patent application EP-A-0 489 968, and as the E1 protein is expected to be subject to immune attack as part of the viral envelope and expected to contain epitopes, the NS4, Core and E1 epitopes of the new types and subtypes disclosed herein will consistently differ from the epitopes present in previously known genotypes. This is examplified by the type-specificity of NS4 synthetic peptides as described in Simmonds et al. (1993c) and Stuyver et al. (1993b) and PCT/EP 94/01323 and the type-specificity of recombinant E1 proteins as described in Maertens et al. (1994).

The peptides according to the present invention contain preferably at least 3, preferably 4, 5 contiguous HCV amino acids, 6, 7 preferably however at least 8 contiguous HCV amino acids, at least 10 or at least 15 (for instance at least 9, 10, 11, 12, 13, 14, 15, 16, 17, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50 or more amino acids).

TABLE 4

30	Amino acids	Synonymous groups
	Ser (S)	Ser, Thr, Gly, Asn
	Arg (R)	Arg, His, Lys, Glu, Gln

		•
	Leu (L)	Leu; Ile, Met, Phe, Val, Tyr
	Pro (P)	Pro, Ala, Thr, Gly
	Thr (T)	Thr, Pro, Ser, Ala, Gly, His, Gln
	Ala (A)	Ala, Pro, Gly, Thr
E	Val (V)	Val, Met, Ile, Tyr, Phe, Leu, Val
5	Gly (G)	Gly, Ala, Thr, Pro, Ser
	lle (I)	lle, Met, Leu, Phe, Val, Ile, Tyr
		Phe, Met, Tyr, Ile, Leu, Trp, Val
	Phe (F)	Tyr, Phe, Trp, Met, Ile, Val, Leu
	Tyr (Y)	Cys, Ser, Thr, Met
10	Cys (C)	2 .
	His (H)	His, Gln, Arg, Lys, Glu, Thr
	GIn (Q)	Gln, Glu, His, Lys, Asn, Thr, Arg
	Asn (N)	Asn, Asp, Ser, Gln
	Lys (K)	Lys, Arg, Glu, Gln, His
15	Asp (D)	Asp, Asn, Glu, Gln
15	· · · · · · · · · · · · · · · · · · ·	Glu, Gln, Asp, Lys, Asn, His, Arg
	Glu (E)	
	Met (M)	Met, Ile, Leu, Phe, Val

<u>Table 4</u> Overview of the amino acid substitutions which could form the basis of analogs (muteins) as defined above

The polypeptides of the invention, and particularly the fragments, can be prepared by classical chemical synthesis.

The synthesis can be carried out in homogeneous solution or in solid phase.

For instance, the synthesis technique in homogeneous solution which can be used is the one described by Houbenweyl in the book entitled "Methode der organischen chemie" (Method of organic chemistry) edited by E. Wunsh, vol. 15-I et II. THIEME, Stuttgart 1974.

The polypeptides of the invention can also be prepared in solid phase according to the methods described by Atherton and Shepard in their book entitled "Solid phase peptide synthesis" (IRL Press, Oxford, 1989).

The polypeptides according to this invention can be prepared by means of recombinant DNA techniques as described by Maniatis et al., Molecular Cloning: A Laboratory Manual, New York, Cold Spring Harbor Laboratory, 1982).

The present invention relates particularly to a polypeptide as defined above, comprising in its amino acid sequence at least one of the following amino acid residues:

20

25

30

10

15

20

25

30

115, C38, V44, A49, Q43, P49, Q55, A58, S60 or D60, E68 or V68, H70, A71 or Q71 or N71, D72, H81, H101, D106, S110, L130, I134, E135, L140, S148, T150 or E150, Q153, F155, D157, G160, E165, I169, F181, L186, T190, T192 or I192 or H192, I193, A195, S196, R197 or N197 or K197, Q199 or D199 or H199, N199, F200 or T200, A208, I213, M216 or S216, N217 or S217 or G217 or K217, T218, I219, A222, Y223, I230, W231 or L231, S232 or H232 or A232, Q233, E235 or L235, F236 or T236, F237, L240 or M240, A242, N244, N249, I250 or K250 or R250, A252 or C252, A254, I255 or V255, D256 or M256, E257, E260 or K260, R261, V268, S272 or R272, I285, G290 or F290, A291, A293 or L293 or W293, T294 or A294, S295, H295, K296 or E296, Y297 or M297, I299 or Y299, I300, S301, P316, S2646, A2648, G2649, A2650, V2652, Q2653, H2656 or L2656, D2657, F2659, K2663 or Q2663, A2667 or V2667, D2677, L2681, M2686 or Q2686 or E2686, A2692 or K2692, H2697, I2707, L2708 or Y2708, A2709, A2719 or M2719, F2727, T2728 or D2728, E2729, F2730 or Y2730, 12741, 12745, V2746 or E2746 or L2746 or K2746, A2748, S2749 or P2749, R2750, E2751, D2752 or N2752 or S2752 or T2752 or V2752 or I2752 or Q2752, S2753 or D2753 or G2753, D2754, A2755, L2756 or Q2756, or R2757,

with said notation being composed of a letter representing the amino acid residue by its one-letter code, and a number representing the amino acid numbering according to Kato et al., 1990 as shown in Table 1 (see also the numbering in Figures 2, 4 and 6),

or a part thereof which is unique to at least one of the HCV subtypes or types as defined in Table 5, and which contains at least one amino acid differing from any of the known HCV types or subtypes, or an analog thereof being substantially homologous and biologically equivalent to said polypeptide or part thereof.

These unique amino acid residues can be deduced from aligning the new HCV amino acid sequences as given in Figure 3 to all known HCV sequences. An alignment with the new sequences as represented in SEQ ID NO 1 to 106 is given in for instance Figures 2, 4 and 6. It should be clear that the alignments given in these figures may be completed with all known HCV sequences to illustrate that any of the above-given unique residues is indeed unique for at least one of the new HCV sequences of the present invention.

Within the group of unique and new amino acid residues of the present

10

15

20

25

30

invention, unique residues may be found which are specific for the following new types (subtypes) of HCV according to the HCV classification system used in the present invention: type 1 subtype 1d, 1e, 1f or 1g isolates; type 2 subtype 2e, 2f, 2g, 2h, 2i, 2k or 2l isolates; type 3 subtype 3g isolates; type 4 subtype 4k, 4l or 4m isolates; type 7 subtype 7a, 7c or 7d isolates, type 9, type 10 or type 11 isolates. In order to obtain these residues the alignments given in Figures 2, 4 and 6 may be used to deduce the type- and or subtype-specificity of any of the unique residues given above.

For example T190 (detected in subtype 1d) refers to a threomine at position 190 (see Figure 2). In other sequences only a serine (S190) or exceptionally an alanine (A190 in type 10a) can be detected.

The polypeptides according to this embodiment of the invention may be possibly labelled, or attached to a solid substrate, or coupled to a carrier molecule such as biotin, or mixed with a proper adjuvant all known in the art and according to the intended use (diagnostic, therapeutic or prophylactic).

The present invention also relates to a polypeptide as defined above, comprising in its amino acid sequence at least one of the sequences repesented by SEQ ID NO107 to 207 as listed above, or a part thereof which is unique to at least one of the HCV subtypes or types as defined in Table 5, or an analog thereof being substantially homologous and biologically equivalent to said polypeptide or part thereof.

The present invention relates also to a polypeptide having an amino acid sequence as represented in any of SEQ ID NO 1 to 106, or a part thereof which is unique to at least one of the HCV subtypes or types as defined in Table 5, or an analog thereof being substantially homologous and biologically equivalent to said polypeptide or part thereof.

The variable region in the core protein (V-CORE in Fig. 2) has been shown to be useful for serotyping (Machida et al., 1992). The sequence of the type 1 subtype 1d, 1e, 1f or 1g sequence; type 2 subtype 2e, 2f, 2g, 2h, 2i, 2k and 2l sequence; type 3 subtype 3g; type 4, subtype 4k, 4l or 4m sequence; type 7 (subtype 7a, 7c and 7d sequences), 9, 10 or 11 sequences of the present invention show type-specific features in this region. The peptide from amino acid 68 to 78 (V-core region) shows the following unique sequence for the sequences of the present invention (see

10

15

20

25

figure	21	٠
nguic	-	

ARQSDGRSWAQ or ARRSEGRSWAQ as for	subtype 1d (SEQ ID NO 107 and
108)	
ERRPEGRSWAQ as for subtype 1e	(SEQ ID NO 109)

ARRPEGRSWAQ as for subtype 1e (SEQ ID NO 109)

ARRPEGRSWAQ as for subtype 1f (SEQ ID NO 110)

DRRTTGKSWGR as for subtype 2k (SEQ ID NO 111)

DRRATGRSWGR as for subtype 2e (SEQ ID NO 112)

DRRATGKSWGR as for subtype 2f (SEQ ID NO 113)

VRQPTGRSWGQ as for type 9 (SEQ ID NO 114)

VRHQTGRTWAQ as for subtype 7a and 7c (SEQ ID NO 115)

VRQNQGRTWAQ as for subtype 7d (SEQ ID NO 116)

ARRTEGRSWAQ as for type 10 (SEQ ID NO 117)

VRRTTGRXXXX or VRRTTGRTWAQ as for type 11

(SEQ ID NO 118 and

119)

Five type-specific variable regions (V1 to V5) can be identified after aligning E1 amino acid sequences of the genotypes of the present invention to the genotypes already known, as shown in Figure 2.

Region V1 encompasses amino acids 192 to 203, this is the amino-terminal 10 amino acids of the E1 protein. The following unique sequences as shown in Fig. 2 can be deduced:

HEVRNASGVYHV or HEVRNASGVYHL as for subtype 1d, (SEQ ID NO 120 and 121)

YEVHSTIDGYHV as for subtype 1f	(SEQ ID NO 122)
VEVKNTSQAYMA as for subtype 2e	(SEQ ID NO 123)
IQVKNNSHFYMA as for subtype 2f	(SEQ ID NO 124)
VQVKNTSTMYMA as for subtype 2g	(SEQ ID NO 125)
VQVKNTSHSYMV as for subtype 2h	(SEQ ID NO 126)
VQVANRSGSYMV as for subtype 2i	(SEQ ID NO 127)

VEIKNTXNTYVL or VEIKNTSNTYVL as for subtype 2k (SEQ ID NO 128

30 and 129)

INYRNVSGIYYV or INYRNTSGIYHV or INYHNTSGIYHI or TNYRNVSGIYHV a for subtype 4k (SEQ ID NO 130, 131, 132 or 133)

QHYRNVSGIYHV as for subtype 4I (SEQ ID NO 134)

	30			
	IQVKNASGIYHL as for type 9	(SEQ ID NO 135)		
•	AHYTNKSGLYHL as for subtype 7c	(SEQ ID NO 136)		
	LNYANKSGLYHL as for subtype 7d	(SEQ ID NO 137)		
	LEYRNASGLYMV as for type 10	(SEQ ID NO 138)		
5	Region V2 encompasses amino acids	s 213 to 223. The following unique		
₹.	sequences can be found in the V2 region as shown in Figure 2:			
٠.	IYEMDGMIMHY or IYEMSGMILHA as for subtype 1d, (SEQ ID NO 139			
	and 140)			
	VYEAKDIILHT as for subtype 1f	(SEQ ID NO 141)		
10	VWQLXDAVLHV as for subtype 2e	(SEQ ID NO 142)		
	VWQLRDAVLHV as for subtype 2f	(SEQ ID NO 143)		
	IWQMQGAVLHV as for subtupe 2g	(SEQ ID NO 144)		
	VWQLKDAVLHV as for subtype 2h	(SEQ ID NO 145)		
	VWQLEEAVLHV as for subtype 2i	(SEQ ID NO 146)		
15	TWQLXXAVLHV as for subtype 2k	(SEQ ID NO 147)		
	VYEADHHILHL or VYEADHHILAL or VFEADHHILHL as for subtype 4k (SEQ ID NO 148, 149 and 150)			
	VYESDHHILHL as for subtype 4I	(SEQ ID NO 151)		
	VFEAETMILHL as for type 9	(SEQ ID NO 152)		
20	VYEAETLILHL as for subtype 7c	(SEQ ID NO 153)		
	VYEANGMILHL as for subtype 7d	(SEQ ID NO 154)		
	VYEAGDIILHL as for type 10.	(SEQ ID NO 155)		
	Region V3 encompasses the amino ac	ids 230 to 242. The following unique		
	V3 region sequences can be deduced from F			
25	VREDNHLRCWMAL or VRENNSSRCW	MAL as for subtype 1d		
	(SEQ ID NO	156 and 157)		
	IREGNISRCWVLP as for subtype 1f	(SEQ ID NO 158)		
	ENSSGRFHCWIPI as for subtype 2e	(SEQ ID NO 159)		
	ERSGNRTFCWTAV as for subtype 2f	(SEQ ID NO 160)		
30	ELQGNKSRCWIPV as for subtype 2g	(SEQ ID NO 162)		
	ERHQNQSRCWIPV as for subtype 2h	(SEQ ID NO 163)		
	EWKDNTSRCWIPV as for subtype 2i	(SEQ ID NO 164)		
•	EREGNSSRCWIPV as for subtype 2k	(SEQ ID NO 165)		

10

15

20

25

30

31 VREGNOSRCWVAL or VRVGNOSSCWVAL or VRVGNQSRCWVAL or VKEGNHSRCWVAL as for subtype 4k (SEQ ID NO 166, 167, 168 or 169) VKTGNTSRCWVAL as for subtype 4I (SEQ ID NO 170) IKAGNESRCWLPV as for type 9 (SEQ ID NO 171) VKXXNQSRCWVQA as for subtype 7c (SEQ ID NO 172) VKTGNLTKCWLSA as for subtype 7d (SEQ ID NO 173) VRSGNTSRCWIPV as for type 10 (SEQ ID NO 174) Region V4 encompasses the amino acids 248 to 257. The following unique V4 region sequences can be deduced from figure 2: VKNASVPTAA or VKDANVPTAA as for subtype 1d (SEQ ID NO 175 and 176) ARIANAPIDE as for subtype 1f (SEQ ID NO 177) VSKPGALTKG as for subtype 2e (SEQ ID NO 178) VSRPGALTRG as for subtype 2f (SEQ ID NO 179) VNQPGALTRG as for subtype 2g (SEQ ID NO 180) VSQPGALTRG as for subtype 2h (SEQ ID NO 181) VSQPGALTKG as for subtype 2i (SEQ ID NO 182) VSRPGALTEG as for subtype 2k (SEQ ID NO 183) APYIGAPLES or APYTAAPLES as for subtype 4k (SEQ ID NO 184 and 185) APILSAPLMS as for subtype 4I (SEQ ID NO 186) VPNSSVPIHG as for type 9 (SEQ ID NO 187) VPNASTPVTG as for subtype 7c. (SEQ ID NO 188) VQNASVSIRG as for subtype 7d (SEQ ID NO 189) VKSPCAATAS as for type 10 (SEQ ID NO 190) Region V5 encompasses the amino acids 294 to 303. The following unique V5 region peptides can be deduced from figure 2: SPRMHHTTQE or SPRLYHTTQE as for subtype 1d (SEQ ID NO 191 and 192) TSRRHWTVQD as for subtype 1f

SUBSTITUTE SHEET (RULE 26)

APKRHYFVQE as for subtype 2e

SPQYHTFVQE as for subtype 2f

SPQHHNFSQD as for subtype 2g

Charles !

.. (SEQ ID NO 193)

(SEQ ID NO 194)

(SEQ ID NO 195)

(SEQ ID NO 196)

10

15

20

25

30

(SEQ ID NO 197) SPQHHIFVQD as for subtype 2h

(SEQ ID NO 198) SPEHHHFVQD as for subtype 2k

RPRRHWTTQD or RPRRHWTAQD or QPRRHWTTQD or RPRRHWTTQE as for (SEQ ID NO 199, 200, 201 or 202) subtype 4k

(SEQ ID NO 203) QPRRHWTVQD as for subtype 41

(SEQ ID NO 204) RPKYHQVTQD as for type 9

(SEQ ID NO 205) RPRMHQVVQE as for subtype 7c

(SEQ ID NO 206) RPRMYEIAQD as for subtype 7d

(SEQ ID NO 207) RHRQHWTVQD as for type 10

The above given list of peptides are particularly useful for treatment and vaccine and diagnostic development.

Also comprised in the present invention is any synthetic peptide (see below) or polypeptide containing at least an epitope derived from the above-defined peptides in their peptidic chain. Also comprised within the present invention is any synthetic peptide or polypeptide comprising at least 6, 7, 8, or 9 contiguous amino acids derived from the above-defined peptides in their peptidic chain.

As used herein, 'epitope' or 'antigenic determinant' means an amino acid sequence that is immunoreactive. Generally an epitope consists of at least 3 to 4 amino acids, and more usually, consists of at least 5 or 6 amino acids, sometimes the epitope consists of about 7 to 8, or even about 10 amino acids.

The present invention particularly relates to any peptide (see below) or polypeptide contained in any of the amino acid sequences as represented in SEQ ID NO 2, 4, 7, 9, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104 or 106 (see Table 5 and Figure 3, Examples section).

The present invention also relates to a recombinant polypeptide encoded by a polynucleic acid as defined above, or a part thereof which is unique to any of the HCV subtypes or types as defined in Table 5, or an analog thereof being substantially homologous and biologically equivalent to said polypeptide.

The present invention also relates to a recombinant expression vector comprising a polynucleic acid or a part thereof as defined above, operably linked to prokaryotic, eukaryotic or viral transcription and translation control elements.

10

15

20

25

30

In general said recombinant vector will comprise a vector sequence, an appropriate prokaryotic, eukaryotic or viral promoter sequence followed by the nucleotide sequences as defined above, with said recombinant vector allowing the expression of any one of the polypeptides as defined above in a prokaryotic, or eukaryotic host or in living mammals when injected as naked DNA, and more particularly a recombinant vector allowing the expression of any of the new HCV sequences of the invention spanning particularly the following amino acid positions:

- a polypeptide starting in the region between positions 1 and 10 and ending at any position in the region between positions 70 and 420, more particularly a polypeptide spanning positions 1 to 70, 1 to 85, positions 1 to 120, positions 1 to 150, positions 1 to 191, or positions 1 to 200, for expression of the Core protein, and a polypeptide spanning positions 1 to 263, positions 1 to 326, positions 1 to 383, or positions 1 to 420 for expression of the Core and E1 protein;
- a polypeptide starting at any position in the region between positions 117 and 192, and ending at any position in the region between positions 263 and 420, for expression of E1, or forms that have the hydrophobic region deleted (positions 264 to 293 plus or minus 8 amino acids);
- a polypeptide starting at any position in the region between positions 1556 and 1688, and ending at any position in the region between positions 1739 and 1764, for expression of NS4, more particularly; a polypeptide starting at position 1658 and ending at position 1711, for expression of NS4a antigen, and more particularly, a polypeptide starting at position 1712 and ending in the region between positions 1743 and 1972 (for instance 1712-1743, 1712-1764, 1712-1782, 1712-1972, 1712-1782, 1712-1902), for expression of NS4b antigen or parts thereof.

Any other HCV vector construction known in the art may also be used for the recombinant polypeptides of the present invention.

Also any of the known purification methods for recombinant proteins may be used for the production of the recombinant polypeptides of the present invention, particularly the HCV recombinant polypeptide purification methods as disclosed in PCT/EP 95/03031 in name of Innogenetics N.V.

The term "vector" may comprise a plasmid, a cosmid, a phage, or a virus or

10

15

20

25

30

a transgenic animal. Particularly useful for vaccine development may be BCG or adenoviral vectors, as well as avipox recombinant viruses.

The present invention also relates to a method for the production of a recombinant polypeptide as defined above, comprising:

transformation of an appropriate cellular host with a recombinant vector, in which a polynucleic acid or a part thereof according to as defined above has been inserted under the control of appropriate regulatory elements,

culturing said transformed cellular host under conditions enabling the expression of said insert, and,

harvesting said polypeptide.

The term 'recombinantly expressed' used within the context of the present invention refers to the fact that the proteins of the present invention are produced by recombinant expression methods be it in prokaryotes, or lower or higher eukaryotes as discussed in detail below.

The term 'lower eukaryote' refers to host cells such as yeast, fungi and the like. Lower eukaryotes are generally (but not necessarily) unicellular. Preferred lower within Saccharomyces, species particularly eukaryotes yeasts, are Schizosaccharomyces, Kluveromyces, Pichia (e.g. Pichia pastoris), Hansenula (e.g. Schizosaccharomyces, Schwaniomyces, Yarowia, Hansenula polymorpha), Zygosaccharomyces and the like. Saccharomyces cerevisiae, S. carlsbergensis and K. lactis are the most commonly used yeast hosts, and are convenient fungal hosts.

The term 'prokaryotes' refers to hosts such as <u>E.coli</u>, <u>Lactobacillus</u>, <u>Lactococcus</u>, <u>Salmonella</u>, <u>Streptococcus</u>, <u>Bacillus subtilis</u> or <u>Streptomyces</u>. Also these hosts are contemplated within the present invention.

The term 'higher eukaryote' refers to host cells derived from higher animals, such as mammals, reptiles, insects, and the like. Presently preferred higher eukaryote host cells are derived from Chinese hamster (e.g. CHO), monkey (e.g. COS and Vero cells), baby hamster kidney (BHK), pig kidney (PK15), rabbit kidney 13 cells (RK13), the human osteosarcoma cell line 143 B, the human cell line HeLa and human hepatoma cell lines like Hep G2, and insect cell lines (e.g. Spodoptera frugiperda). The host cells may be provided in suspension or flask cultures, tissue cultures, organ cultures and the like. Alternatively the host cells may also be transgenic animals.

The term 'recombinant polynucleotide or nucleic acid' intends a polynucleotide

10

15

20

25

30

or nucleic acid of genomic, cDNA, semisynthetic, or synthetic origin which, by virtue of its origin or manipulation: (1) is not associated with all or a portion of a polynucleotide with which it is associated in nature, (2) is linked to a polynucleotide other than that to which it is linked in nature, or (3) does not occur in nature.

The term 'recombinant host cells', 'host cells', 'cells', 'cell lines', 'cell cultures', and other such terms denoting microorganisms or higher eukaryotic cell lines cultured as unicellular entities refer to cells which can be or have been, used as recipients for a recombinant vector or other transfer polynucleotide, and include the progeny of the original cell which has been transfected. It is understood that the progeny of a single parental cell may not necessarily be completely identical in morphology or in genomic or total DNA complement as the original parent, due to natural, accidental, or deliberate mutation.

The term 'replicon' is any genetic element, e.g., a plasmid, a chromosome, a virus, a cosmid, etc., that behaves as an autonomous unit of polynucleotide replication within a cell; i.e., capable of replication under its own control.

The term 'vector' is a replicon further comprising sequences providing replication and/or expression of a desired open reading frame.

The term 'control sequence' refers to polynucleotide sequences which are necessary to effect the expression of coding sequences to which they are ligated. The nature of such control sequences differs depending upon the host organism; in prokaryotes, such control sequences generally include promoter, ribosomal binding site, splicing sites and terminators; in eukaryotes, generally, such control sequences include promoters, splicing sites, terminators and, in some instances, enhancers. The term 'control sequences' is intended to include, at a minimum, all components whose presence is necessary for expression, and may also include additional components whose presence is advantageous, for example, leader sequences which govern secretion.

The term 'promoter' is a nucleotide sequence which is comprised of consensus sequences which allow the binding of RNA polymerase to the DNA template in a manner such that mRNA production initiates at the normal transcription initiation site for the adjacent structural gene.

The expression 'operably linked' refers to a juxtaposition wherein the components so described are in a relationship permitting them to function in their

10

15

20

25

30

intended manner. A control sequence 'operably linked' to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences.

The segment of the HCV cDNA encoding the desired sequence inserted into the vector sequence may be attached to a signal sequence. Said signal sequence may be that from a non-HCV source, e.g. the IgG or tissue plasminogen activator (tpa) leader sequence for expression in mammalian cells, or the α -mating factor sequence for expression into yeast cells, but particularly preferred constructs according to the present invention contain signal sequences appearing in the HCV genome before the respective start points of the proteins.

A variety of vectors may be used to obtain recombinant expression of HCV single or specific oligomeric envelope proteins of the present invention. Lower eukaryotes such as yeasts and glycosylation mutant strains are typically transformed with plasmids, or are transformed with a recombinant virus. The vectors may replicate within the host independently, or may integrate into the host cell genome.

Higher eukaryotes may be transformed with vectors, or may be infected with a recombinant virus, for example a recombinant vaccinia virus. Techniques and vectors for the insertion of foreign DNA into vaccinia virus are well known in the art, and utilize, for example homologous recombination. A wide variety of viral promoter sequences, possibly terminator sequences and poly(A)-addition sequences, possibly enhancer sequences and possibly amplification sequences, all required for the mammalian expression, are available in the art. Vaccinia is particularly preferred since vaccinia halts the expression of host cell proteins. Vaccinia is also very much preferred since it allows the expression of f.i. E1 and E2 proteins of HCV in cells or individuals which are immunized with the live recombinant vaccinia virus. For vaccination of humans the avipox and Ankara Modified Virus (AMV) are particularly useful vectors.

Also known are insect expression transfer vectors derived from baculovirus Autographa californica nuclear polyhedrosis virus (AcNPV), which is a helper-independent viral expression vector. Expression vectors derived from this system usually use the strong viral polyhedrin gene promoter to drive the expression of heterologous genes. Different vectors as well as methods for the introduction of the heterologous DNA into the desired site of baculovirus are available to the man skilled

in the art for baculovirus expression. Also different signals for posttranslational modification recognized by insect cells are known in the art.

The present invention also relates to a host cell transformed with a recombinant vector as defined above.

5

10

The present invention also relates to a method for detecting antibodies to HCV present in a biological sample, comprising:

- (i) contacting the biological sample to be analysed for the presence of HCV with a polypeptide as defined above,
- (ii) detecting the immunological complex formed between said antibodies and said polypeptide.

The present invention also relates to a method for HCV typing, comprising:

- (i) contacting the biological sample to be analysed for the presence of HCV with a polypeptide as defined above,
- (ii) detecting the immunological complex formed between said antibodies and said polypeptide.

The present invention also relates to a diagnostic kit for use in detecting the presence of HCV, said kit comprising at least one polypeptide as defined above, with said polypeptide being preferably bound to a solid support.

20

15

The present invention also relates to a diagnostic kit for HCV typing, said kit comprising at least one polypeptide as defined above, with said polypeptide being preferably bound to a solid support.

The present invention also relates to diagnostic kit according as defined above, said kit comprising a range of said polypeptides which are attached to specific locations on a solid substrate.

25

30

The present invention also relates to a diagnostic kit as defined above, wherein said solid support is a membrane strip and said polypeptides are coupled to the membrane in the form of parallel lines.

The immunoassay methods according to the present invention may utilize antigens from the different domains of the new and unique polypeptide sequences of the present invention that maintain linear (in case of peptides) and conformational epitopes (in case of polypeptides) recognized by antibodies in the sera from individuals infected with HCV. It is within the scope of the invention to use for instance single or specific oligomeric antigens, dimeric antigens, as well as

10

15

20

25

30

combinations of single or specific oligomeric antigens. The HCVantigens of the present invention may be employed in virtually any assay format that employs a known antigen to detect antibodies. Of course, a format that denatures the HCV conformational epitope should be avoided or adapted. A common feature of all of these assays is that the antigen is contacted with the body component suspected of containing HCV antibodies under conditions that permit the antigen to bind to any such antibody present in the component. Such conditions will typically be physiologic temperature, pH and ionic strenght using an excess of antigen. The incubation of the antigen with the specimen is followed by detection of immune complexes comprised of the antigen.

Design of the immunoassays is subject to a great deal of variation, and many formats are known in the art. Protocols may, for example, use solid supports, or immunoprecipitation. Most assays involve the use of labeled antibody or polypeptide; the labels may be, for example, enzymatic, fluorescent, chemiluminescent, radioactive, or dye molecules. Assays which amplify the signals from the immune complex are also known; examples of which are assays which utilize biotin and avidin or streptavidin, and enzyme-labeled and mediated immunoassays, such as ELISA assays.

The immunoassay may be, without limitation, in a heterogeneous or in a homogeneous format, and of a standard or competitive type. In a heterogeneous format, the polypeptide is typically bound to a solid matrix or support to facilitate separation of the sample from the polypeptide after incubation. Examples of solid supports that can be used are nitrocellulose (e.g., in membrane or microtiter well form), polyvinyl chloride (e.g., in sheets or microtiter wells), polystyrene latex (e.g., in beads or microtiter plates, polyvinylidine fluoride (known as ImmunolonTM), diazotized paper, nylon membranes, activated beads, and Protein A beads. For example, Dynatech ImmunolonTM 1 or ImmunlonTM 2 microtiter plates or 0.25 inch polystyrene beads (Precision Plastic Ball) can be used in the heterogeneous format. The solid support containing the antigenic polypeptides is typically washed after separating it from the test sample, and prior to detection of bound antibodies. Both standard and competitive formats are know in the art.

In a homogeneous format, the test sample is incubated with the combination, of antigens in solution. For example, it may be under conditions that will precipitate

any antigen-antibody complexes which are formed. Both standard and competitive formats for these assays are known in the art.

In a standard format, the amount of HCV antibodies in the antibody-antigen complexes is directly monitored. This may be accomplished by determining whether labeled anti-xenogeneic (e.g. anti-human) antibodies which recognize an epitope on anti-HCV antibodies will bind due to complex formation. In a competitive format, the amount of HCV antibodies in the sample is deduced by monitoring the competitive effect on the binding of a known amount of labeled antibody (or other competing ligand) in the complex.

10

5

Complexes formed comprising anti-HCV antibody (or in the case of competitive assays, the amount of competing antibody) are detected by any of a number of known techniques, depending on the format. For example, unlabeled HCV antibodies in the complex may be detected using a conjugate of anti-xenogeneic lg complexed with a label (e.g. an enzyme label).

15

In an immunoprecipitation or agglutination assay format the reaction between the HCV antigens and the antibody forms a network that precipitates from the solution or suspension and forms a visible layer or film of precipitate. If no anti-HCV antibody is present in the test specimen, no visible precipitate is formed.

20

There currently exist three specific types of particle agglutination (PA) assays. These assays are used for the detection of antibodies to various antigens when coated to a support. One type of this assay is the hemagglutination assay using red blood cells (RBCs) that are sensitized by passively adsorbing antigen (or antibody) to the RBC. The addition of specific antigen antibodies present in the body component, if any, causes the RBCs coated with the purified antigen to agglutinate.

25

To eliminate potential non-specific reactions in the hemagglutination assay, two artificial carriers may be used instead of RBC in the PA. The most common of these are latex particles. However, gelatin particles may also be used. The assays utilizing either of these carriers are based on passive agglutination of the particles coated with purified antigens.

30 .

The HCV antigens of the present invention comprised of conformational epitopes will typically be packaged in the form of a kit for use in these immunoassays. The kit will normally contain in separate containers the native HCV antigen, control antibody formulations (positive and/or negative), labeled antibody



10

15

20

25

30

when the assay format requires the same and signal generating reagents (e.g. enzyme substrate) if the label does not generate a signal directly. The native HCV antigen may be already bound to a solid matrix or separate with reagents for binding it to the matrix. Instructions (e.g. written, tape, CD-ROM, etc.) for carrying out the assay usually will be included in the kit.

Immunoassays that utilize the native HCV antigen are useful in screening blood for the preparation of a supply from which potentially infective HCV is lacking. The method for the preparation of the blood supply comprises the following steps. Reacting a body component, preferably blood or a blood component, from the individual donating blood with HCV polypeptides of the present invention to allow an immunological reaction between HCV antibodies, if any, and the HCV antigen. Detecting whether anti-HCV antibody - HCV antigen complexes are formed as a result of the reacting. Blood contributed to the blood supply is from donors that do not exhibit antibodies to the native HCV antigens.

In cases of a positive reactivity to the HCV antigen, it is preferable to repeat the immunoassay to lessen the possibility of false positives. For example, in the large scale screening of blood for the production of blood products (e.g. blood transfusion, plasma, Factor VIII, immunoglobulin, etc.) 'screening' tests are typically formatted to increase sensitivity (to insure no contaminated blood passes) at the expense of specificity; i.e. the false-positive rate is increased. Thus, it is typical to only defer for further testing those donors who are 'repeatedly reactive'; i.e. positive in two or more runs of the immunoassay on the donated sample. However, for confirmation of HCV-positivity, the 'confirmation' tests are typically formatted to increase specificity (to insure that no false-positive samples are confirmed) at the expense of sensitivity.

The solid phase selected can include polymeric or glass beads, nitrocellulose, microparticles, microwells of a reaction tray, test tubes and magnetic beads. The signal generating compound can include an enzyme, a luminescent compound, a chromogen, a radioactive element and a chemiluminescent compound. Examples of enzymes include alkaline phosphatase, horseradish peroxidase and betagalactosidase. Examples of enhancer compounds include biotin, anti-biotin and avidin. Examples of enhancer compounds binding members include biotin, anti-biotin and avidin. In order to block the effects of rheumatoid factor-like substances, the

10

15

20

25

30

test sample is subjected to conditions sufficient to block the effect of rheumatoid factor-like substances. These conditions comprise contacting the test sample with a quantity of anti-human lgG to form a mixture, and incubating the mixture for a time and under conditions sufficient to form a reaction mixture product substantially free of rheumatoid factor-like substance.

The present invention particularly relates to an immunoassay format in which the polypeptides (or peptides) of the invention are coupled to a membrane in the form of parallel lines. This assay format is particularly advantageous for HCV typing purposes.

The present invention also relates to a pharmaceutical composition comprising at least one (recombinant) polypeptides as defined above and a suitable excipient, diluent or carrier.

The present invention also relates to a method of preventing HCV infection, comprising administering the pharmaceutical composition as defined above to a mammal in effective amount to stimulate the production of protective antibody or protective T-cell response.

The present invention relates to the use of a composition as defined above in a method for preventing HCV infection.

The present invention further relates to a vaccine for immunizing a mammal against HCV infection, comprising at least one (recombinant) polypeptide as defined above, in a pharmaceutically acceptable carrier.

The term 'immunogenic' refers to the ability of a substance to cause a humoral and/or cellular response, whether alone or when linked to a carrier, in the presence or absence of an adjuvant. 'Neutralization' refers to an immune response that blocks the infectivity, either partially or fully, of an infectious agent. A 'vaccine' is an immunogenic composition capable of eliciting protection against HCV, whether partial or complete. A vaccine may also be useful for treatment of an individual, in which case it is called a therapeutic vaccine.

The term 'therapeutic' refers to a composition capable of treating HCV infection. The term 'effective amount' refers to an amount of epitope-bearing polypeptide sufficient to induce an immunogenic response in the individual to which it is administered, or to otherwise detectably immunoreact in its intended system (e.g., immunoassay). Preferably, the effective amount is sufficient to effect

10

15

20

25

30

treatment, as defined above. The exact amount necessary will vary according to the application. For vaccine applications or for the generation of polyclonal antiserum / antibodies, for example, the effective amount may vary depending on the species, age, and general condition of the individual, the severity of the condition being treated, the particular polypeptide selected and its mode of administration, etc. It is also believed that effective amounts will be found within a relatively large, non-critical range. An appropriate effective amount can be readily determined using only routine experimentation. Preferred ranges of proteins for prophylaxis of HCV disease are 0.01 to 100 μ g/dose, preferably 0.1 to 50 μ g/dose. Several doses may be needed per individual in order to achieve a sufficient immune response and subsequent protection against HCV disease.

The present invention also relates to a vaccine as defined above, comprising at least one (recombinant) polypeptide as defined above, with said polypeptide being unique for at least one of the subtypes or types as defined above.

Said vaccine compositions may include prophylactic as well as therapeutic vaccine compositions.

Pharmaceutically acceptable carriers include any carrier that does not itself induce the production of antibodies harmful to the individual receiving the composition. Suitable carriers are typically large, slowly metabolized macromolecules such as proteins, polysaccharides, polylactic acids, polyglycolic acids, polymeric amino acids, amino acid copolymers; and inactive virus particles. Such carriers are well known to those of ordinary skill in the art.

Preferred adjuvants to enhance effectiveness of the composition include, but are not limited to: aluminim hydroxide (alum), N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP) as found in U.S. Patent No. 4,606,918, N-acetyl-normuramyl-L-alanyl-D-isoglutamine (nor-MDP), N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-sn-glycero-3-hydroxyphosphoryloxy)-ethylamine (MTP-PE) and RIBI, which contains three components extracted from bacteria, monophosphoryl lipid A, trehalose dimycolate, and cell wall skeleton (MPL+TDM+CWS) in a 2% squalene/Tween 80 emulsion. Any of the 3 components MPL, TDM or CWS may also be used alone or combined 2 by 2. Additionally, adjuvants such as Stimulon (Cambridge Bioscience, Worcester, MA)

10

15

20

25

30

Immunogenic compositions used as vaccines comprise a 'sufficient amount' or 'an immunologically effective amount' of the proteins of the present invention, as well as any other of the above mentioned components, as needed. 'Immunologically effective amount', means that the administration of that amount to an individual, either in a single dose or as part of a series, is effective for treatment, as defined above. This amount varies depending upon the health and physical condition of the individual to be treated, the taxonomic group of individual to be treated (e.g. nonhuman primate, primate, etc.), the capacity of the individual's immune system to synthesize antibodies, the degree of protection desired, the formulation of the vaccine, the treating doctor's assessment of the medical situation, the strain of infecting HCV, and other relevant factors. It is expected that the amount will fall in a relatively broad range that can be determined through routine trials. Usually, the amount will vary from 0.01 to 1000 μ g/dose, more particularly from 0.1 to 100 μ g/dose.

The proteins of the invention may also serve as vaccine carriers to present homologous (e.g. T cell epitopes or B cell epitopes fromfor istance the core,E1, E2, NS2, NS3, NS4 or NS5 regions) or heterologous (non-HCV) haptens, in the same manner as Hepatitis B surface antigen (see European Patent Application 174,444). In this use, envelope proteins provide an immunogenic carrier capable of stimulating an immune response to haptens or antigens conjugated to the aggregate. The antigen may be conjugated either by conventional chemical methods, or may be cloned into the gene encoding E1 and/or E2 at a location corresponding to a hydrophilic region



10

15

20

25

30

of the protein. Such hydrophylic regions include the V1 region (encompassing amino acid positions 191 to 202), the V2 region (encompassing amino acid positions 213 to 223), the V3 region (encompassing amino acid positions 230 to 242), the V4 region (encompassing amino acid positions 230 to 242), the V5 region (encompassing amino acid positions 294 to 303) and the V6 region (encompassing amino acid positions 329 to 336). Another useful location for insertion of haptens is the hydrophobic region (encompassing approximately amino acid positions 264 to 293). It is shown in the present invention that this region can be deleted without affecting the reactivity of the deleted E1 protein with antisera. Therefore, haptens may be inserted at the site of the deletion.

The immunogenic compositions are conventionally administered parenterally, typically by injection, for example, subcutaneously or intramuscularly. Additional formulations suitable for other methods of administration include oral formulations and suppositories. Dosage treatment may be a single dose schedule or a multiple dose schedule. The vaccine may be administered in conjunction with other immunoregulatory agents.

The administration of the immunogen(s) of the present invention may be for either a prophylactic or therapeutic purpose. When provided prophylactically, the immunogen(s) is provided in advance of any exposure to HCV or in advance of any symptom of any symptoms due to HCV infection. The prophylactic administration of the immunogen serves to prevent or attenuate any subsequent infection of HCV in a mammal. When provided therapeutically, the immunogen(s) is provided at (or shortly after) the onset of the infection or at the onset of any symptom of infection or disease caused by HCV. The therapeutic administration of the immunogen(s) serves to attenuate the infection or disease.

In addition to use as a vaccine, the compositions can be used to prepare antibodies to HCV (E1) proteins. The antibodies can be used directly as antiviral agents. To prepare antibodies, a host animal is immunized using the E1 proteins native to the virus particle bound to a carrier as described above for vaccines. The host serum or plasma is collected following an appropriate time interval to provide a composition comprising antibodies reactive with the (E1) protein of the virus particle. The gamma globulin fraction or the IgG antibodies can be obtained, for example, by use of saturated ammonium sulfate or DEAE Sephadex, or other

10

15

20

25

30

techniques known to those skilled in the art. The antibodies are substantially free of many of the adverse side effects which may be associated with other anti-viral agents such as drugs.

The present invention also relates particularly to a peptide corresponding to an amino acid sequence encoded by at least one of the HCV genomic sequences as defined above, with said peptide being unique to any of the HCV subtypes or types as defined in Table 5, and which contains at least one amino acid differing from any of the known HCV types or subtypes, or an analog thereof being substantially homologous and biologically equivalent.

The present invention relates particularly to a peptide comprising at least one unique epitope of the new sequences of the invention as represented in SEQ ID NO 1 to 106.

The present invention relates also particularly to a peptide comprising in its sequence a unique amino acid residue of the invention as defined above.

The present invention relates particularly to a peptide which is biotinylated as explained in WO 93/18054.

All the embodiments (immunoassay formats, vaccines, compositions, uses, etc.) illustrated for the polypeptides of the invention as above also relate to the peptides of the invention.

The present invention also relates to a method for detecting antibodies to HCV present in a biological sample, comprising:

- (i) contacting the biological sample to be analysed for the presence of HCV with a peptide as defined above,
- (ii) detecting the immunological ccomplex formed between said antibodies and said peptide.

The present invention also relates to a method for HCV typing, comprising: (i) contacting the biological sample to be analysed for the presence of HCV with a peptide as defined above,

(ii) detecting the immunological complex formed between said antibodies and said peptide.

The present invention also relates to a diagnostic kit for use in detecting the presence of HCV, said kit comprising at least one peptide as defined above, with said peptide being preferably bound to a solid support.

The present invention also relates to a diagnostic kit for HCV typing, said kit comprising at least one peptide as defined above, with said peptide being preferably bound to a solid support.

The present invention also relates to a diagnostic kit as defined above, wherein said peptides are selected from the following:

- at least one NS4 peptide,
- at least one NS4 peptide and at least one Core peptide,
- at least one NS4 peptide and at least one Core peptide and at least one E1 peptide,
- at least one NS4 peptide and at least one E1 peptide.

The present invention also relates to a diagnostic kit as defined above, said kit comprising a range of said peptides which are attached to specific locations on a solid substrate.

The present invention also relates to a diagnostic kit as defined above, wherein said solid support is a membrane strip and said peptides are coupled to the membrane in the form of parallel lines.

The present invention also relates to a pharmaceutical composition comprising at least one as defined above and a suitable excipient, diluent or carrier.

the present invention also relates to a method of preventing HCV infection, comprising administering the pharmaceutical composition as defined above to a mammal in effective amount to stimulate the production of protective antibody or protective T-cell response.

The present invention also relates to the use of a composition as defined above in a method for preventing HCV infection.

The present invention also relates to a vaccine for immunizing a mammal against HCV infection, comprising at least one peptide as defined above, in a pharmaceutically acceptable carrier.

The present invention relates also to a vaccine as defined above, comprising at least one peptide as defined above, with said peptide being unique for at least one of the subtypes or types as defined in Table 5.

The present invention relates to an antibody raised upon immunization with at least one polypeptide or peptide as defined above, with said antibody being specifically reactive with any of said polypeptides or peptides, and with said antibody being preferably a monoclonal antibody.

-

15

10

5

20

25

30

The monoclonal antibodies of the invention can be produced by any hybridoma liable to be formed according to classical methods from splenic cells of an animal, particularly from a mouse or rat, immunized against the HCV polypeptides according to the invention as defined above on the one hand, and of cells of a myeloma cell line on the other hand, and to be selected by the ability of the hybridoma to produce the monoclonal antibodies recognizing the polypeptides which has been initially used for the immunization of the animals.

The antibodies involved in the invention can be labelled by an appropriate label of the enzymatic, fluorescent, or radioactive type.

10

5

The monoclonal antibodies according to this preferred embodiment of the invention may be humanized versions of mouse monoclonal antibodies made by means of recombinant DNA technology, departing from parts of mouse and/or human genomic DNA sequences coding for H and L chains or from cDNA clones coding for H and L chains.

15

20

Alternatively the monoclonal antibodies according to this preferred embodiment of the invention may be human monoclonal antibodies. These antibodies according to the present embodiment of the invention can also be derived from human peripheral blood lymphocytes of patients infected with HCV type 1 subtype 1d, 1e, 1f or 1g, HCV type 2 subtype 2e, 2f, 2g, 2h, 2i, 2k or 2l; HCV type 3, subtype 3g; HCV type 4 subtype 4k, 4l or 4m; and/or HCV type 7 (subtypes 7a, 7c or 7d), 9, 10 or 11, or vaccinated against HCV. Such human monoclonal antibodies are prepared, for instance, by means of human peripheral blood lymphocytes (PBL) repopulation of severe combined immune deficiency (SCID) mice (for recent review, see Duchosal et al. 1992) or by screening Eppstein Barr-virustransformed lymphocytes of infected or vaccinated individuals for the presence of reactive B-cells by means of the antigens of the present invention.

25

The invention also relates to the use of the proteins of the invention, muteins thereof, or peptides derived therefrom for the selection of recombinant antibodies by the process of repertoire cloning (Persson et al., 1991).

30

Antibodies directed to peptides derived from a certain genotype may be used either for the detection of such HCV genotypes, or as therapeutic agents.

The present invention relates also to a method for detecting HCV antigens present in a biological sample, comprising:

10

15

20

25

30

- (i) contacting said biological sample with an antibody as defined above,
- (ii) detecting the immune compleexes formed between said HCV antigens and said antibody.

The present invention relates also to a method for HCV typing, comprising:

- (i) contacting said biological sample with an antibody as defined above,
- (ii) detecting the immune complexes formed between said HCV antigens and said antibody.

The present invention relates also to a diagnostic kit for use in detecting the presence of HCV, said kit comprising at least one antibody as defined above, with said antibody being preferably bound to a solid support.

The present invention relates also to a diagnostic kit for HCV typing, said kit comprising at least one antibody as defined above, with said antibody being preferably bound to a solid support.

The present invention relates also to a diagnostic kit as defined above, said kit comprising a range of said antibodies which are attached to specific locations on a solid substrate.

The present invention relates also to a pharmaceutical composition comprising at least one antibody as defined above and a suitable excipient, diluent or carrier.

The present invention relates also to a method of preventing or treating HCV infection, comprising administering the pharmaceutical composition as defined above to a mammal in effective amount.

The present invention relates also to the use of a composition as defined above in a method for preventing or treating HCV infection.

The genotype may also be detected by means of a type-specific antibody as defined above, which may also linked to any polynucleotide sequence that can afterwards be amplified by PCR to detect the immune complex formed (Immuno-PCR, Sano et al., 1992).

Any publications or patent applications referred to herein are incorporated by reference. The following examples illustrate aspects of the invention but are in no way intended to limit the scope thereof.

FIGURE LEGENDS

Figure Legends

Figure 1

Alignment of the nucleotide sequences of the Core/E1 region of some of the isolates of the newly identified types and subtypes of the present invention, with other known prototype isolates of subtypes.

Figure 2

5

10

Alignment of the amino acid sequences of the Core/E1 region of some of the isolates of the newly identified types and subtypes of the present invention, with other known prototype isolates of subtypes.

Figure 3

Nucleotide and amino acid sequences obtained from the new HCV isolates of the present invention (SEQ ID NO 1 to 106).

Figure 4

Alignment of the amino acid sequences of the Core/E1 region of some of the isolates of the newly identified types and subtypes of the present invention, with other known prototype isolates of subtypes.

Figure 5

Alignment of the nucleotide sequences of the NS5b region of some of the isolates of the newly identified types and subtypes of the present invention, with other known prototype isolates of subtypes.

Figure 6

Alignment of the amino acid sequences of the NS5b region of some of the isolates of the newly identified types and subtypes of the present invention, with other known prototype isolates of subtypes.

5 Table 5

Overview of the new subtypes and types of the present invention and the regions sequenced. The subtypes between barckets have been replaced by the non-bracketed subtypes following the classification of Tokita et al. (1994).

Examples

10 Serum samples.

15

25

Serum samples from Cameroonian blood donors (CAM) were screened for HCV antibodies with Innotest HCV Ab III, and confirmed by INNO-LIA HCV III (Innogenetics, Antwerp, Belgium). Serum samples from patients with chronic hepatitis C infection were obtained from various centers in the Benelux countries (BNL), from France (FR), from Pakistan (PAK), from Egypt (EG), and from Vietnam (VN).

Samples from the Benelux, Cameroon, France and Vietnam were selected because of their aberrant reactivities (isolates CAM1078, FR2, FR1, VN4, VN12, VN13, NE98 and others (see Table 5)).

20 cPCR, LiPA, cloning and sequencing.

RNA isolation, cDNA synthesis, PCR, cloning, and LiPA genotyping using biotinylated 5' UR amplification products were performed as described (Stuyver et al., 1994c). The 5' UR, the Core/E1, and the NS5B PCR products were used for direct sequencing. The sequence of the universal 5' UR primers HCPr95, HCPr96, HCPr98, and HCPr29, were described previously (Stuyver et al. 1993b). The following primers were also described (Stuyver et al. 1994c): HCPr41, a sense

10

15

20

primer for the amplification of the Core region; HCPr52 and HCPr54 for amplification of the Core/E1 region; and HCPr206 and HCPr207 for amplication of a 340-bp NS5B region.

Serum samples BNL1, BNL2, BNL3, BNL4, BNL5, BNL6, BNL7, BNL8, BNL9, BNL10, BNL11, BNL12, CAM1078, FR2, FR16, FR4, FR13, VN13, VN4, VN12, FR1, NE98, and FR19 were analyzed in the Core/E1 region by direct sequencing. Serum samples BNL1, BNL2, FR17, CAM1078, FR2, FR16, BNL3, FR4, BNL5, FR13, FR18, PAK64, BNL8, BNL12, EG81, VN13, VN4, VN12, FR1, NE98, FR14, FR15, and FR19 were also analyzed in the NS5B region by direct sequencing. Partial 5' UR, Core, E1, and NS5B sequences were obtained. The length of the obtained sequences is sufficient to classify the obtained sequences into new types or subtypes, based on the phylogenetic distances to known sequences. The following sequences could be obtained (nucleotide sequences have odd-numbered SEQ ID NO., amino acid sequences have even-numbered SEQ ID NO.): SEQ ID NO 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87, 89, 91, 93, 95, 97, 99, 101, 103 and 105. The amino acid sequences deduced therefrom are given in SEQ ID NO 2, 4, 7, 9, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104 and 106. Table 5 gives an overview of these sequences.

5/13590	Ė	PCT/EP95/04155
0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	7932-8271(SEQ ID NO. 71) 7932-8271(SEQ ID NO. 73) 7932-8271(SEQ ID NO. 77) ¹³ 7932-8271(SEQ ID NO. 79) 7932-8271(SEQ ID NO. 81) 7932-8271 (SEQ ID NO. 83)	7932-8271 (SEQ ID NO. 85) 7932-8271 (SEQ ID NO. 87) 7932-8271 (SEQ ID NO. 89) 7932-8271 (SEQ ID NO. 91) 7932-8271 (SEQ ID NO. 93) 7932-8271 (SEQ ID NO. 95) 7932-8271 (SEQ ID NO. 95) 7932-8271 (SEQ ID NO. 97) 7932-8266 (SEQ ID NO. 99) 7932-8271 (SEQ ID NO. 99)
	**	
A78-925 (SEQ ID NO. 3) 478-925 (SEQ ID NO. 7) (-238)-414 (SEQ ID NO. 59) 65) 478-957 (SEQ ID NO. 15)	478-925 (SEQ ID NO. 19) 478-925 (SEQ ID NO. 23) 478-833 (SEQ ID NO. 25) 7.75) 478-925 (SEQ ID NO. 29) 478-925 (SEQ ID NO. 31) 478-925 (SEQ ID NO. 33) 478-925 (SEQ ID NO. 33) 478-925 (SEQ ID NO. 33)	-925 (SEQ ID NO.
¬ — — — — —	1-957 (SEQ ID NO 17) 1-310 (SEQ ID NO. 21) (-238)-957 (SEQ ID NO 17)	1-413 (SEQ ID NO. 45) 1-957 (SEQ ID NO. 43) 1-957 (SEQ ID NO. 47) 1-957 (SEQ ID NO. 41) 1-310 (SEQ ID NO. 49) 478 (-238)-223 (SEQ ID NO. 103)
Table 5 Type Isolate 1d BNL1 1d FR17 1e CAM1078 1f FR2 1g FR16 2e BNL3	2f FR4 2g BNL4 2h BNL5 2i BNL6 2k FR13 2l FR18 3g PAK64 4k BNL7 4k BNL8 4k BNL9 4k BNL9	(8b) (8a) (9a) (7a) a

W C 30/13330	53	PCT/EP95/04155
(SEQ ID NO. (SEQ I	2645-2757 (SEQ ID NO. 72, 2645-2757 (SEQ ID NO. 74) 2645-2757 (SEQ ID NO. 78) 2645-2757 (SEQ ID NO. 80) 2645-2757 (SEQ ID NO. 82) 2645-2757 (SEQ ID NO. 82)	2645-2757 (SEQ ID NO. 86) 2645-2757 (SEQ ID NO. 88) 2645-2757 (SEQ ID NO. 90) 2645-2757 (SEQ ID NO. 92) 2645-2757 (SEQ ID NO. 94) 2645-2757 (SEQ ID NO. 94) 2645-2757 (SEQ ID NO. 98) 2645-2757 (SEQ ID NO. 100) 2645-2757 (SEQ ID NO. 102) 2645-2757 (SEQ ID NO. 102)
sequence position 159-308 (SEQ ID NO. 4) 159-308 (SEQ ID NO. 8) 1-138 (SEQ ID NO. 60) 159-317 (SEQ ID NO. 16)	159-308 (SEQ ID NO. 20) 159-308 (SEQ ID NO. 24) 159-277 (SEQ ID NO. 26) 159-308 (SEQ ID NO. 30) 159-308 (SEQ ID NO. 32) 159-308 (SEQ ID NO. 34) 159-308 (SEQ ID NO. 34)	ID NO.
Amino acid 1-103 (SEQ ID NO. 2) 1-103 (SEQ ID NO. 6) 1-74 (SEQ ID NO. 10) 1-316 (SEQ ID NO. 12) 1-103 (SEQ ID NO. 66) 1-317 (SEQ ID NO. 14)	N N ON ON	1-137 (SEQ ID NO. 46) 1-317 (SEQ ID NO. 44) 1-317 (SEQ ID NO. 42) 1-103 (SEQ ID NO. 50) 1-74 (SEQ ID NO. 104)
pe	29 BNL4 2h BNL5 2i BNL6 2k FR13 21 FR18 39 PAK64 4k BNL7 4k BNL9 4k BNL9 4k BNL9	(8b) (9a) (7a)

Phylogenetic analysis.

Previously published sequences were taken from the EMBL/Genbank database. Alignments were created using the program HCVALIGN (Stuyver et al. 1994c). Sequences were presented in a sequential format to the Phylogeny Inference Package (PHYLIP) version 3.5c (public domain program freely available from the University of Washington, Seattle, USA). Distance matrices were produced by DNADIST using the Kimura 2-parameter setting and further analyzed in NEIGHBOR, using the neighborioining setting. The program DRAWTREE was used to create graphic outputs.

Identification of new subtypes

10

15

5

These analyses indicated the clustering of BNL1, BNL2, CAM 1078, FR2, FR16, and FR17 with type 1 isolates, yet neither of these sequences clustered together with any of the known type 1 subtypes 1a, 1b, or 1c. BNL1, BNL2, and FR17 clearly clustered together and could be assigned a new type 1 subtype 1d, while CAM1078 could be classified into another new subtype 1e, FR2 could be classified into another type 1 subtype 1f, and FR16 could be classified into yet another type 1 subtype 1g. Interestingly, all 3 type 1d isolates (BNL1, BNL2, and FR17) and 1g isolate FR16 were obtained from patients of Moroccan ethnic origin who resided in Europe.

.

20

25

Another group of isolates showed homology to other type 2 sequences, but none of the isolates BNL3, FR4, BNL4, BNL5, BNL6, FR13, or FR18 could be classified into one of the known type 2 subtypes 2a, 2b, 2c (Bukh et al., 1993), or 2d (Stuyver et al., 1994c). Based on the phylogenetic distances to other type 2 isolates and to other isolates of the group, each of these isolates could be classified into a new type 2 subtype. BNL3 was assigned subtype 2e, FR4 subtype 2f, BNL4 subtype 2g, BNL5 subtype 2h, and BNL6 could be classified into yet another type 2 subtype 2i. If the previously published isolate HN4 is classified as 2j, FR13 and FR18 may be classified into new type 2 subtypes 2k and 2l. However, the possibility that FR13 and FR18 could belong to subtypes 2g or 2i has not yet been ruled out. Definite classification can be obtained by determining the NS5B sequences of isolates BNL4 and BNL6, belonging to subtypes 2g and 2i, respectively.

30

Isolate PAK64 showed homology to type 3 sequences, but could not be classified into one of the known type 3 subtypes 3a to f. Based on the phylogenetic distances to other type 3 isolates, PAK64 could be classified into a new type 3

10

subtype. PAK64 was assigned subtype 3g. However, the possibility that PAK64 belongs to a known type 3 subtype can not be strictly ruled out since only one region of the genome has been sequenced. Definite classification can be obtained by determining the Core/E1 sequences of isolate PAK64 after amplification with primerHcPr52 and HcPr54.

Among the Benelux and Egyptian samples that were analyzed, some sequences clustered with the previously identified type 4 subtypes 4c and 4d. However, BNL7, BNL8, BNL9, BNL10, BNL11, BNL12, and EG81 clustered into new subtypes of type 4. Isolates BNL7, BNL8, BNL9, BNL10, and BNL11 clustered again separately from BNL12 and EG81 into a new subtype 4k. This subtype was the predominant subtype in the Benelux countries. BNL12 and EG81 also segregated into separate subtypes. BNL12 was assigned to another new subtype 4l and EG81 was assigned to yet another new subtype 4m.

10

15

Identification of new HCV major types

Isolates FR1, VN4, VN12, VN13, NE98, FR14, FR15, and FR19 did not cluster with any of the known 6 major types of HCV. VN4, VN12, and VN13 were very distantly related to genotype 6, but phylogenetic analysis indicated that these isolates should be assigned new major types. VN13, VN4 and VN12 were related at the subtype level and assigned type 7a, 7c, and 7d, respectively. FR1 was not related to any known isolate and was assigned genotype 9a. NE98 shows a distant relatedness to type 3 sequences, yet phylogenetic analysis suggested classification into a new major type 10a. Depending on international guidelines for assigning type and subtype levels, NE98 may also be classified into an additional type 3 subtype. FR14, FR15, and FR19 show a very distant relatedness to type 2 sequences, yet phylogenetic analysis indicated thes isolates to be classified into a new major type 11, all belonging to the same subtype designated 11a. Depending on international guidelines for assigning type and subtype levels, FR14, FR15, and FR19 may also be classified into an additional type 2 subtype.

REFERENCES

Barany F (1991). Genetic disease detection and DNA amplification using cloned thermostable ligase. Proc Natl Acad Sci USA 88: 189-193.

Bej A, Mahbubani M, Miller R, Di Cesare J, Haff L, Atlas R (1990) Mutiplex PCR amplification and immobilized capture probes for detection of bacterial pathogens and indicators in water. Mol Cell Probes 4:353-365.

Bukh J, Purcell R, Miller R (1992). Sequence analysis of the 5' noncoding region of hepatitis C virus. Proc Natl Acad Sci USA 89:4942-4946.

Bukh J, Purcell R, Miller R (1993). At least 12 genotypes of hepatitis C virus predicted by sequence analysis of the putative E1 gene of isolates collected worldwide. Proc. Natl. Acad. Sci. USA 90,8234-8238.

Cha T, Beal E, Irvine B, Kolberg J, Chien D, Kuo G, Urdea M (1992) At least five related, but distinct, hepatitis C viral genotypes exist. Proc Natl Acad Sci USA 89:7144-7148.

15 Chan S-W, Simmonds P, McOmish F, Yap P, Mitchell R, Dow B, Follett E (1991) Serological responses to infection with three different types of hepatitis C virus. Lancet 338:1991.

Chan S-W, McOmish F, Holmes E, Dow B, Peutherer J, Follett E, Yap P, Simmonds P (1992) Analysis of a new hepatitis C virus type and its phylogenetic relationship to existing variants. J Gen Virol 73:1131-1141.

Chomczynski P, Sacchi N (1987) Single step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. Anal Biochem 162:156-159.

Choo Q, Richman K, Han J, Berger K, Lee C, Dong C, Gallegos C, Coit D, Medina-SUBSTITUTE SHEET (RULE 26)

Selby A, Barr P, Weiner A, Bradley D, Kuo G, Houghton M (1991) Genetic organization and diversity of the hepatitis C virus. Proc Natl Acad Sci USA 88:2451-2455.

Compton J (1991). Nucleic acid sequence-based amplification. Nature, 350: 91-92.

Duchosal A, Eming S, Fisher P (1992) Immunization of hu-PBL-SCID mice and the resue of human monoclonal Fab fragments through combinatorial libraries. Nature 355:258-262.

Duck P (1990). Probe amplifier system based on chimeric cycling oligonucleotides. Biotechniques 9, 142-147.

Guatelli J, Whitfield K, Kwoh D, Barringer K, Richman D, Gengeras T (1990)

10 Isothermal, in vitro amplification of nucleic acids by a multienzyme reaction modeled after retroviral replication. Proc Natl Acad Sci USA 87: 1874-1878.

Hijikata M, Kato N, Ootsuyama Y, Nakagawa M, Shimotohmo K (1991) Gene mapping of the putative structural region of the hepatitis C virus genome by in vitro processing analysis. Proc Natl Acad Sci USA 88, 5547-5551.

Jacobs K, Rudersdorf R, Neill S, Dougherty J, Brown E, Fritsch E (1988) The thermal stability of oligonucleotide duplexes is sequence independent in tetraalkylammonium salt solutions: application to identifying recombinant DNA clones. Nucl Acids Res 16:4637-4650.

Kato N, Hijikata M, Ootsuyama Y, Nakagawa M, Ohkoshi S, Sugimura T, Shimotohno K (1990) Molecular cloning of the human hepatitis C virus genome from Japanese patients with non-A, non-B hepatitis. Proc Natl Acad Sci USA 87:9524-9528.

Kwoh D, Davis G, Whitfield K, Chappelle H, Dimichele L, Gingeras T (1989). Transcription-based amplification system and detection of amplified human immunodeficiency virus type 1 with a bead-based sandwich hybridization format. Proc Natl Acad Sci USA, 86: 1173-1177.

20

Kwok S, Kellogg D, McKinney N, Spasic D, Goda L, Levenson C, Sinisky J, (1990). Effects of primer-template mismatches on the polymerase chain reaction: Human immunodeficiency views type 1 model studies. Nucl. Acids Res., 18: 999.

Landgren U, Kaiser R, Sanders J, Hood L (1988). A ligase-mediated gene detection technique. Science 241:1077-1080.

Lizardi P, Guerra C, Lomeli H, Tussie-Luna I, Kramer F (1988) Exponential amplification of recombinant RNA hybridization probes. Bio/Technology 6:1197-1202.

Lomeli H, Tyagi S, Printchard C, Lisardi P, Kramer F (1989) Quantitative assays based on the use of replicatable hybridization probes. Clin Chem 35: 1826-1831.

Machida A, Ohnuma H, Tsuda F, Munekata E, Tanaka T, Akahane Y, Okamoto H, Mishiro S (1992) Hepatology 16, 886-891.

Maniatis T, Fritsch E, Sambrook J (1982) Molecular cloning: a laboratory manual. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.

Mori S, Kato N, Yagyu A, Tanaka T, Ikeda Y, Petchclai B, Chiewsilp P, Kurimura T, Shimotohno K (1992) A new type of hepatitis C virus in patients in Thailand. Biochem Biophys Res Comm 183:334-342.

Okamoto H, Okada S, Sugiyama Y, Kurai K, lizuka H, Machida A, Miyakawa Y, Mayumi M (1991) Nucleotide sequence of the genomic RNA of hepatitis C virus isolated from a human carrier: comparison with reported isolates for conserved and divergent regions. J Gen Virol 72:2697-2704.

Okamoto H, Kurai K, Okada S, Yamamoto K, Lizuka H, Tanaka T, Fukuda S, Tsuda F, Mishiro S (1992) Full-length sequences of a hepatitis C virus genome having poor homology to reported isolates: comparative study of four distinct genotypes. Virology 188:331-341.

20

Persson M, Caothien R, Burton D (1991). Generation of diverse high-affinity human monoclonal antibodies by repertoire cloning. Proc Natl Acad Sci USA 89:2432-2436. Saiki R, Gelfand D, Stoffel S, Scharf S, Higuchi R, Horn G, Mullis K, Erlich H (1988). Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. Science 239:487-491.

Saiki R, Walsh P, Levenson C, Erlich H (1989) Genetic analysis of amplified DNA with immobilized sequence-specific oligonucleotide probes (1989) Proc Natl Acad Sci USA 86:6230-6234.

Sano T, Smith C, Cantor C (1992) Immuno-PCR: very sensitive antigen detection by means of specific antibody-DNA conjugates. Science 258:120-122.

Simmonds P, McOmsh F, Yap P, Chan S, Lin C, Dusheiko G, Saeed A, Holmes E (1993a), Sequence variability in the 5' non-coding region of hepatitis C virus: identification of a new virus type and restrictions on sequence diversity. J Gen Virology, 74:661-668.

Stuyver L, Rossau R, Wyseur A, Duhamel M, Vanderborght B, Van Heuverswyn H, Maertens G (1993b) Typing of hepatitis C virus (HCV) isolates and characterization of new (sub)types using a Line Probe Assay. J Gen Virology, 74: 1093-1102.

Tokita et al. (1994) Hepatitis C virus vraiants from Vietnam are classifiable into the seventh, eighth, and ninth major genetic groups. Proc. Natl. Acad. Sci, 91: 11022-11026.

Walker G, Little M, Nadeau J, Shank D (1992). Isothermal in vitro amplification of DNA by a restriction enzyme/DNA polymerase system. Proc Natl Acad Sci USA 89:392-396.

Wu D, Wallace B (1989). The ligation amplification reaction (LAR) - amplification of specific DNA sequences using sequential rounds of template-dependent ligation. Genomics 4:560-569.

15

20

Miller P, Yano J, Yano E, Carroll C, Jayaram K, Ts'o P (1979) Nonionic nucleic acid analogues. Synthesis and characterization of dideoxyribonucleoside methylphosphonates. Biochemistry 18(23):5134-43.

Nielsen P, Egholm M, Berg R, Buchardt O (1991) Sequence-selective recognition of DNA by strand displacement with a thymine-substituted polyamide. Science 254(5037):1497-500.

Nielsen P, Egholm M, Berg R, Buchardt O (1993) Sequence specific inhibition of DNA restriction enzyme cleavage by PNA. Nucleic-Acids-Res. 21(2):197-200.

Asseline U, Delarue M, Lancelot G, Toulme F, Thuong N (1984) Nucleic acid-binding molecules with high affinity and base sequence specificity: intercalating agents covalently linked to oligodeoxynucleotides. Proc. Natl. Acad. Sci. USA 81(11):3297-301.

Matsukura M, Shinozuka K, Zon G, Mitsuya H, Reitz M, Cohen J, Broder S (1987) Phosphorothioate analogs of oligodeoxynucleotides: inhibitors of replication and cytopathic effects of human immunodeficiency virus. Proc. Natl. Acad. Sci. USA 84(21):7706-10.

Maertens, G., Ducatteeuw, A., Stuyver, L., Vandeponseele, P., Venneman, A., Wyseur, A., Bosman, F., Heijtink, R. & de Martynoff, G. (1994) Low prevalence of anti-E1 antibodies reactive to recombinant type 1b E1 envelope protein in type 2, 3, and 4 sera, but high prevalence in subtypes 1a and 1b. In: Viral Hepatitis and Liver Disease, Proceedings of the International Symposium on Viral Hepatitis and Liver Disease (Eds. Nishioka, K., Suzuki, H., Mishiro, S., and Oda, T.), pp 314-316, Springer-Verlag Tokyo.

Simmonds, P., Rose, K.A., Graham, S., Chan, S.-W., McOmish, F., Dow, B.C., Follett, E.A.C., Yap, P.L., & Marsden, H. (1993b) Mapping of serotype-specific, immunodominant epitopes in the NS4 region of hepatitis C virus (HCV): Use of type-specific peptides to serologically discriminate infections with HCV type 1, 2, and 3.

J. Clin. Microbiol. 31, 1493-1503.

Simmonds, P., Holmes, E.C., Cha, T.-A., Chan, S.-W., McOmish, F., Irvine, B., Beall, E., Yap, P.L., Kolberg, J., & Urdea, M.S. (1993c) *J. Gen. Virol.* **74**, 2391-2399.

Stuyver, L., Van Arnhem, W., Wyseur, A. & Maertens, G. (1994) Cloning and phylogenetic analysis of the Core, E2, and NS3/4 regions of hepatitis C virus type 5a. Biochem. Biophys. Res. Comm. 202, 1308-1314.

Simmonds, P., Alberti, A., Alter, H., Bonino, F., Bradley, D.W., Bréchot, C., Brouwer, J., Chan, S.-W., Chayama K., Chen, D.-S., Choo, Q.-L., Colombo, M., Cuypers, T., Date, T., Dusheiko, G., Esteban, J.I., Fay, O., Hadziyannis, S., Han, J., Hatzakis, A.,
Holmes, E.C., Hotta, H., Houghton, M., Irvine, B., Kohara, M., Kolberg, J.A., Kuo, G., Lau, J.Y.N., Lelie, P.N., Maertens, G., McOmish, F., Miyamura, T., Mizokami, M., Nomoto, A., Prince A.M., Reesink, H.W., Rice, C., Roggendorf, M., Schalm, S., Shikata, T., Shimotohno, K., Stuyver, L., Trépo, C., Weiner, A., Yap, P.L. & Urdea, M.S. (1994) A proposed system for the nomenclature of hepatitis C virus genotypes.
Hepatology 19, 1321-1324.

Stuyver, L., Van Arnhem, W., Wyseur, A., DeLeys, R. & Maertens, G. (1993a) Analysis of the putative E1 envelope and NS4a epitope regions of HCV type 3. Biochem. Biophys. Res. Comm. **192**, 635-641.

Stuyver, L., Rossau, R., Wyseur, A., Duhamel, M., Vanderborght, B., Van Heuverswyn, H. & Maertens, G. (1993b) Typing of hepatitis C virus isolates and characterization of new subtypes using a line probe assay. J. Gen Virol. **74**, 1093-1102.

Stuyver, L., Wyseur, A., Van Arnhem, W., Rossau, R., Delaporte, E., Dazza, M.-C., Van Doorn, L.-J., Kleter, B. & Maertens, G. (1994a) The use of a line probe assay as a tool to detect new types or subtypes of hepatitis C virus. In: Viral Hepatitis and Liver Disease, Proceedings of the International Symposium on Viral Hepatitis and Liver Disease (Eds. Nishioka, K., Suzuki, H., Mishiro, S., and Oda, T.), pp 317-319,

Springer-Verlag Tokyo.

Stuyver, L., Van Arnhem, W., Wyseur, A. & Maertens, G. (1994b) Cloning and Phylogenetic analysis of the Core, E2, and NS3/4 regions of the hepatitis C virus type 5a. Biochem. Biophys. Res. Comm. 202, 1308-1314.

- Stuyver, L., Van Arnhem, W., Wyseur, A., Hernandez, F., Delaporte, E., & Maertens, G. (1994c) Classification of hepatitis C viruses based on phylogenetics analysis of the E1 and NS5B regions and identification of 5 new subtypes. Proc. Natl. Acad. Sci. USA 91.
- Stuyver et al. (1995) Hepatitis C virus genotyping by means of 5'-UR/core line probe assays and molecular analysis of untypeable samples. Virus Reasearch (in press).

15

CLAIMS

- 1. An HCV polynucleic acid, having a nucleotide sequence which is unique to a theretofore unidentified HCV type or subtype which is different from HCV subtypes 1a, 1b, 1c, 2a, 2b, 2c, 2d, 3a, 3b, 3c, 3d, 3e, 3f, 4a, 4b, 4c, 4d, 4e, 4f, 4g, 4h, 4i, 4j, 5a or 6a, with said HCV subtypes being classified as in Table 3 by comparison of a part of the NS5 gene nucleotide sequence spanning positions 7932 to 8271, with said amino acid numbering being shown in Table 1, and with said polynucleic acid containing at least one nucleotide differing from said known HCV nucleotide sequences, or the complement thereof.
- 2. A polynucleic acid according to claim 1, having a nucleotide sequence which is unique to at least one of HCV subtypes 1d, 1e, 1f, 1g, 2e, 2f, 2g, 2h, 2i, 2k, 2l, 3g, 4k, 4l, 4m, 7a, 7c or 7d, with said HCV subtypes being classified as defined in claim 1.
 - 3. A polynucleic acid according to claim 1, having a nucleotide sequence which is unique to at least one of HCV types 9, 10 or 11, with said HCV types being classified as defined in claim 1.
 - 4. A polynucleic acid according to any of claims 1 to 3 encoding an HCV polyprotein comprising in its amino acid sequence at least one of the following amino acid residues:
- 115, C38, V44, A49, Q43, P49, Q55, A58, S60 or D60, E68 or V68, H70, A71 or Q71 or N71, D72, H81, H101, D106, S110, L130, l134, E135, L140, S148, T150 or E150, Q153, F155, D157, G160, E165, l169, F181, L186, T190, T192 or l192 or H192, l193, A195, S196, R197 or N197 or K197, Q199 or D199 or H199 or N199, F200 or T200, A208, l213, M216 or S216, N217 or S217 or G217 or K217, T218, l219, A222, Y223, l230, W231 or L231, S232 or H232 or A232, Q233, E235 or L235, F236 or T236, F237, L240 or M240, A242, N244, N249, l250 or K250 or R250, A252 or C252, A254, l255 or V255, D256 or M256, E257, E260 or K260, R261, V268, S272 or R272, l285, G290 or F290, A291, A293 or L293 or W293, T294 or A294, S295 or H295, K296 or E296, Y297 or M297, l299 or Y299, l300,

10

30

S301, P316, S2646, A2648, G2649, A2650, V2652, Q2653, H2656 or L2656, D2657, F2659, K2663 or Q2663, A2667 or V1667, D2677, L2681, M2686 or Q2686 or E2686, A2692 or K2692, H2697, I2707, L2708 or Y2708, A2709, A2719 or M2719, F2727, T2728 or D2728, E2729, F2730 or Y2730, I2741, I2745, V2746 or E2746 or L2746 or K2746, A2748, S2749 or P2749, R2750, E2751, D2752 or N2752 or S2752 or T2752 or V2752 or I2752 or Q2752, S2753 or D2753 or G2753, D2754, A2755, L2756 or Q2756, R2757.

with said notation being composed of a letter representing the amino acid residue by its one-letter code, and a number representing the amino acid numbering as shown in Table 1,

or a part of said polynucleic acid which is unique to at least one of the HCV subtypes or types as defined in claims 2 to 3, and which contains at least one nucleotide differing from known HCV nucleotide sequences, or the complement thereof.

5. A polynucleic acid according to any of claims 1 to 4, with said polynucleic acid
 encoding a HCV polyprotein comprising in its amino acid sequence at least one amino acid sequence chosen from the following list:

	3	
	ARQSDGRSWAQ or ARRSEGRSWAQ as for subtype 1d	(SEQ ID NO 107 and 108)
	ERRPEGRSWAQ as for subtype 1e	(SEQ ID NO 109)
	ARRPEGRSWAQ as for subtype 1f	(SEQ ID NO 110)
20	DRRTTGKSWGR as for subtype 2k	(SEQ ID NO 111)
	DRRATGRSWGR as for subtype 2e	(SEQ ID NO 112)
	DRRATGKSWGR as for subtype 2f	(SEQ ID NO 113)
25	VRQPTGRSWGQ as for type 9	(SEQ ID NO 114)
	VRHQTGRTWAQ as for subtype 7a and 7c	(SEQ ID NO 115)
	VRQNQGRTWAQ as for subtype 7d	(SEQ ID NO 116)
	ARRTEGRSWAQ as for type 10	(SEQ ID NO 117)
	VRRTTGRXXXX or VRRTTGRTWAQ as for type 11 119)	(SEQ ID NO 118 and
	HEVRNASGVYHV or HEVRNASGVVHI as for subtract	

HEVRNASGVYHV or HEVRNASGVYHL as for subtype 1d (SEQ ID NO 120 and 121)

YEVHSTTDGYHV as for subtype 1f (SEQ ID NO 122)

VEVKNTSQAYMA as for subtype 2e (SEQ ID NO 123)

IQVKNNSHFYMA as for subtype 2f (SEQ ID NO 124)

	VQVKNTSTMYMA as for subtype 2g	(SEQ ID NO 125)
	VQVKNTSHSYMV as for subtype 2h	(SEQ ID NO 126)
	VQVANRSGSYMV as for subtype 2i	(SEQ ID NO 127)
	VEIKNTXNTYVL or VEIKNTSNTYVL as for	subtype 2k (SEQ ID NO 128 and 129)
5	INYRNVSGIYYV or INYRNTSGIYHV or INY	
	subtype 4k	(SEQ ID NO 130, 131, 132 or 133)
	QHYRNVSGIYHV as for subtype 4I	(SEQ ID NO 134)
	IQVKNASGIYHL as for type 9	(SEQ ID NO 135)
	AHYTNKSGLYHL as for subtype 7c	(SEQ ID NO 136)
10	LNYANKSGLYHL as for subtype 7d	(SEQ ID NO 137)
	LEYRNASGLYMV as for type 10	(SEQ ID NO 138)
	IYEMDGMIMHY or IYEMSGMILHA as for s	ubtype 1d (SEQ ID NO 139 and 140)
	VYEAKDIILHT as for subtype 1f	(SEQ ID NO 141)
	VWQLXDAVLHV as for subtype 2e	(SEQ ID NO 142)
15	VWQLRDAVLHV as for subtype 2f	(SEQ ID NO 143)
-	IWQMQGAVLHV as for subtype 2g	(SEQ ID NO 144)
	VWQLKDAVLHV as for subtype 2h	(SEQ ID NO 145)
	VWQLEEAVLHV as for subtype 2i	(SEQ ID NO 146)
	TWQLXXAVLHV as for subtype 2k	(SEQ ID NO 147)
20	VYEADHHILHL or VYEADHHILAL or VFEA	DHHILHL as for subtupe 4k
	(SE	Q ID NO 148, 149 and 150)
	VYESDHHILHL as for subtype 4I	(SEQ ID NO 151)
	VFEAETMILHL as for type 9	(SEQ ID NO 152)
	VYEAETLILHL as for subtype 7c	(SEQ ID NO 153)
25	VYEANGMILHL as for subtype 7d	(SEQ ID NO 154)
	VYEAGDIILHL as for type 10	(SEQ ID NO 155)
	VREDNHLRCWMAL or VRENNSSRCWMA	L as for subtype 1d
	•	(SEQ ID NO 156 and 157)
	IREGNISRCWVPL as for subtype 1f	(SEQ ID NO 158)
30	ENSSGRFHCWIPI as for subtype 2e	(SEQ ID NO 159)
	ERSGNRTFCWTAV as for subtype 2f	(SEQ ID NO 160)
	ELQGNKSRCWIPV as for subtype 2g	(SEQ ID NO 162)
	ERHQNQSRCWIPV as for subtype 2h	(SEQ ID NO 163)

	·	
	EWKDNTSRCWIPV as for subtype 2i	(SEQ ID NO 164)
	EREGNSSRCWIPV as for subtype 2k	(SEQ ID NO 165)
	VREGNOSRCWVAL or VRTGNOSRCWVAL of	or VRVGNQSSCWVAL or
	VRVGNQSRCWVAL or VKEGNHSRCWVAL as for sub	
5	(SEQ ID NO 16	66, 167, 168 or 169)
	VKTGNTSRCWVAL as for subtype 4I	(SEQ ID NO 170)
	IKAGNESRCWLPV as for type 9	(SEQ ID NO 171)
	VKEGNQSRCWVQA as for subtype 7c	(SEQ ID NO 172)
	VKXXNLTKCWLSA as for subtype 7d	(SEQ ID NO 173)
10	VRSGNTSRCWIPV as for type 10	(SEQ ID NO 174)
	VKNASVPTAA or VKDANVPTAA as for subtype 1d	(SEQ ID NO 175 and
	176)	
	ARIANAPIDE as for subtype 1f	(SEQ ID NO 177)
	VSKPGALTKG as for subtype 2e	(SEQ ID NO 178)
15	VSRPGALTRG as for subtype 2f	(SEQ ID NO 179)
	VNQPGALTRG as for subtype 2g	(SEQ ID NO 180)
	VSQPGALTRG as for subtype 2h	(SEQ ID NO 181)
	VSQPGALTKG as for subtype 2i	(SEQ ID NO 182)
	VSRPGALTEG as for subtype 2k	(SEQ ID NO 183)
. 20	APYIGAPLES or APYTAAPLES as for subtype 4k	(SEQ ID NO 184 and 185)
	APILSAPLMS as for subtype 4I	(SEQ ID NO 186)
	VPNSSVPIHG as for type 9	(SEQ ID NO 187)
	VPNASTPVTG as for subtype 7c	(SEQ ID NO 188)
	VQNASVSIRG as for subtype 7d	(SEQ ID NO 189)
25	VKSPCAATAS as for type 10	(SEQ ID NO 190)
	SPRMHHTTQE or SPRLYHTTQE as for subtype 1d	(SEQ ID NO 191 and 192)
	TSRRHWTVQD as for subtype 1f	(SEQ ID NO 193)
	APKRHYFVQE as for subtype 2e	(SEQ ID NO 194)
	SPQYHTFVQE as for subtype 2f	(SEQ ID NO 195)
30	SPQHHNFSQD as for subtype 2g	(SEQ ID NO 196)
	SPQHHIFVQD as for subtype 2h	(SEQ ID NO 197)
	SPEHHHFVQD as for subtype 2k	(SEQ ID NO 198)
	RPRRHWTTQD or RPRRHWTAQD or QPRRHWTTQD or	RPRRHWTTQE as for

SUBSTITUTE SHEET (RULE 26)

20

25

68

Subtype 4k (SEQ ID NO 199, 200, 201 or 202)

QPRRHWTVQD as for subtype 4l (SEQ ID NO 203)

RPKYHQVTQD as for type 9 (SEQ ID NO 204)

RPRMHQVVQE as for subtype 7c (SEQ ID NO 205)

RPRMYEIAQD as for subtype 7d (SEQ ID NO 206)

RHRQHWTVQD as for type 10 (SEQ ID NO 207)

or a part of said polynucleic acid which is unique to at least one of the HCV subtypes or types as defined in claims 2 to 3, and which contains at least one nucleotide differing from known HCV nucleotide sequences, or the complement thereof.

- 6. A polynucleic acid according to any of claims 1 to 5 having a sequence selected from any of SEQ ID NO 1 to 105, or a part of said polynucleic acid which is unique to at least one of the HCV subtypes or types as defined in claims 2 to 3, and which contains at least one nucleotide differing from known HCV nucleotide sequences, or the complement thereof.
- 7. A polynucleic acid according to any of claims 1 to 6, which codes for the 5' UR, the Core/E1, the NS4 or the NS5B region or a part thereof.
 - 8. A polynucleic acid according to any of claims 1 to 7 which is a cDNA sequence.
 - 9. An oligonucleotide primer comprising part of a polynucleic acid according to any of claims 1 to 8, with said primer being able to act as primer for specifically amplifying the nucleic acid of a certain isolate belonging to the genotype from which the primer is derived.
 - 10. An oligonucleotide probe comprising part of a polynucleic acid according to any of claims 1 to 8, with said probe being able to act as a hybridization probe for specific detection and/or classification into types and/or subtypes of a HCV nucleic acid containing said nucleotide sequence, with said probe being possibly labelled or attached to a solid substrate.
 - 11. A diagnostic kit for use in determing the genotype of HCV, said kit comprising a

primer according to claim 9.

- 12. A diagnostic kit for use in determining the genotype of HCV, said kit comprising a probe according to claim 10.
- 13. A diagnostic kit according to claim 12, wherein said probe(s) is(are) attached to a solid substrate.
- 14. A diagnostic kit according to claim 13, wherein a range of said probes are attached to specific locations on a solid substrate.
- 15. A diagnostic kit according to claim 14, wherein said solid support is a membrane strip and said probes are coupled to the membrane in the form of parallel lines.
- 16. A method for the detection of HCV nucleic acids present in a biological sample, comprising:
 - (i) possibly extracting sample nucleic acid,
 - (ii) amplifying the nucleic acid with at least one primer according to claim 9,
 - (iii) detecting the amplified nucleic acids.
- 15 17. A method for the detection of HCV nucleic acids present in a biological sample, comprising:
 - (i) possibly extracting sample nucleic acid,
 - (ii) possibly amplifying the nucleic acid with at least one primer according to claim 9, or with a universal HCV primer,
- 20 (iii) hybridizing the nucleic acids of the biological sample, possibly under denatured conditions, at appropriate conditions with one or more probes according to claim 10, with said probes being possibly attached to a solid substrate,
 - (iv) possibly washing at appropriate conditions,
- 25 (v) detecting the hybrids formed.
 - 18. A method for detecting the presence of one or more HCV genotypes present in SUBSTITUTE SHEET (RULE 26)

1.5

a biological sample, comprising:

- (i) possibly extracting sample nucleic acid,
- (ii) specifically amplifying the nucleic acid with at least one primer according to claim 9,
- 5 (iii) detecting said amplified nucleic acids,
 - (iv) inferring the presence of one or more genotypes of HCV present from the observed pattern of amplified fragments.
 - 19. A method for detecting the presence of one or more HCV genotypes present in a biological sample, comprising:
- 10 (i) possibly extracting sample nucleic acid,
 - (ii) possibly amplifying the nucleic acid with at least one primer according to claim 9 or with a universal HCV primer,
 - hybridizing the nucleic acids of the biological sample, possibly under denatured conditions, at appropriate conditions with one or more probes according to claim 10, with said probes being possibly attached to a solid substrate,
 - (iv) possibly washing at appropriate conditions,
 - (v) detecting the hybrids formed,
- (vi) inferring the presence of one or more HCV genotypes present from the observed hybridization pattern.
 - 20. A method according to claim 19, wherein said probes are further characterized as defined in any of claims 13 to 15.
 - 21. A method according to claims 16 to 18, wherein said nucleic acids are labelled during or after amplification.
- 22. A polypeptide having an amino acid sequence encoded by a polynucleic acid according to any of claims 1 to 8, or a part thereof which is unique to at least one of the HCV subtypes or types as defined in claims 2 or 3, and which contains at least one amino acid differing from any of the known HCV types or subtypes amino acid sequences, or an analog thereof being substantially homologous and biologically

equivalent.

10

15

20

30

23. A polypeptide according to claim 22 comprising in its amino acid sequence at least one of the following amino acid residues:

115, C38, V44, A49, Q43, P49, Q55, A58, S60 or D60, E68 or V68, H70, A71 or Q71 or N71, D72, H81, H101, D106, S110, L130, I134, E135, L140, S148, T150 5 or E150, Q153, F155, D157, G160, E165, I169, F181, L186, T190, T192 or I192 or H192, I193, A195, S196, R197 or N197 or K197, Q199 or D199 or H199 or N199, F200 or T200, A208, I213, M216 or S216, N217 or S217 or G217 or K217, T218, I219, A222, Y223, I230, W231 or L231, S232 or H232 or A232, Q233, E235 or L235, F236 or T236, F237, L240 or M240, A242, N244, N249, I250 or K250 or R250, A252 or C252, A254, I255 or V255, D256 or M256, E257, E260 or K260, R261, V268, S272 or R272, I285, G290 or F290, A291, A293 or L293 or W293, T294 or A294, S295 or H295, K296 or E296, Y297 or M297, I299 or Y299, I300, S301, P316, S2646, A2648, G2649, A2650, V2652, Q2653, H2656 or L2656, D2657, F2659, K2663 or Q2663, A2667 or V2667, D2677, L2681, M2686 or Q2686 or E2686, A2692 or K2692, H2697, I2707, L2708 or Y2708, A2709, A2719 or M2719, F2727, T2728 or D2728, E2729, F2730 or Y2730, I2741, I2745, V2746 or E2746 or L2746 or K2746, A2748, S2749 or P2749, R2750, E2751, D2752 or N2752 or S2752 or T2752 or V2752 or I2752 or Q2752, S2753 or D2753 or G2753, D2754, A2755, L2756 or Q2756, or R2757,

with said notation being composed of a letter representing the amino acid residue by its one-letter code, and a number representing the amino acid numbering as shown in Table 1,

or a part of said polypeptide which is unique to at least one of the HCV subtypes or types as defined in claims 2 to 3, and which contains at least one amino acid differing from known HCV types or subtypes amino acid sequences, or an analog thereof being substantially homologous and biologically equivalent to said polypeptide.

24. A polypeptide according to claim 22 comprising in its amino acid sequence at least one of the sequences represented by SEQ ID NO 107 to 207 as listed in claim 5, or part of said polypeptide which is unique to at least one of the HCV subtypes or types as defined in claims 2 to 3, and which contains at least one amino acid differing from SUBSTITUTE SHEET (RULE 26)

5

10

15

25

known HCV types or subtypes amino acid sequences, or an analog thereof being substantially homologous and biologically equivalent to said polypeptide.

- 25. A polypeptide having an amino acid sequence as represented in any of SEQ ID NO 1 to 106, or a part thereof which is unique to at least one of the HCV subtypes or types as defined in claims 2 to 3, and which contains at least one amino acid differing from known HCV types or subtypes amino acid sequences, or an analog thereof being substantially homologous and biologically equivalent to said polypeptide.
- 26. A recombinant polypeptide encoded by a polynucleic acid according to any of claims 1 to 8, or a part thereof which is unique to at least one of the HCV subtypes or types as defined in claims 2 or 3, and which contains at least one amino acid differing from known HCV types or subtypes amino acid sequences, or an analog thereof being substantially homologous and biologically equivalent to said polypeptide.
 - 27. A method for production of a recombinant polypeptide of claim 26, comprising:
 - transformation of an appropriate cellular host with a recombinant vector, in which a polynucleic acid or a part thereof according to any of claims 1 to 8 has been inserted under the control of the appropriate regulatory elements,
 - culturing said transformed cellular host under conditions enabling the expression of said insert, and,
 - harvesting said polypeptide.
- 28. A recombinant expression vector comprising a polynucleic acid or a part thereof according to any of claims 1 to 8 operably linked to prokaryotic, eukaryotic or viral transcription and translation control elements.
 - 29. A host cell transformed with a recombinant vector according to claim 28.
 - 30. A method for detecting antibodies to HCV present in a biological sample, comprising:
 - (i) contacting the biological sample to be analysed for the presence of HCV with a polypeptide according to any of claims 22 to 26,

SUBSTITUTE SHEET (RULE 26)

5

10

- (ii) detecting the immunological complex formed between said antibodies and said polypeptide.
- 31. A method for HCV typing, comprising:
- (i) contacting the biological sample to be analysed for the presence of HCV with a polypeptide according to any of claims 22 to 26,
- (ii) detecting the immunological complex formed between said antibodies and said polypeptide.
- 32. A diagnostic kit for use in detecting the presence of HCV, said kit comprising at least one polypeptide according to any of claims 22 to 26, with said polypeptide being possibly bound to a solid support.
- 33. A diagnostic kit for HCV typing, said kit comprising at least one polypeptide according to any of claims 22 to 26, with said polypeptide being possibly bound to a solid support.
- 34. A diagnostic kit according to claims 32 to 33, said kit comprising a range of polypeptides which are attached to specific locations on a solid substrate.
 - 35. A diagnostic kit according to claims 32 to 34, wherein said solid support is a membrane strip and said polypeptides are coupled to the membrane in the form of parallel lines.
- 36. A pharmaceutical composition comprising at least one polypeptide according to
 any of claims 22 to 26 and a suitable excipient, diluent or carrier.
 - 37. A method of preventing HCV infection, comprising administering the pharmaceutical compositon of claim 36 to a mammal in effective amount to stimulate the production of protective antibody or protective T-cell response.
- 38. Use of a composition according to claim 36 in a method for preventing HCV infection as defined in claim 37.

5

10

15

20

25

- 39. A vaccine for immunizing a mammal against HCV infection, comprising at least one polypeptide according to claims 22 to 26, in a pharmaceutically acceptable carrier.
- 40. A vaccine according to claim 39, comprising at least one polypeptide according to claims 22 to 26, with said polypeptide being unique for at least one of the HCV subtypes as defined in claims 2 or 3.
- 41. A peptide corresponding to an amino acid sequence encoded by at least one of the HCV polynucleic acids according to any of claims 1 to 8, with said peptide comprising an epitope being unique to at least one of the HCV subtypes or types as defined in claims 2 or 3, and with said peptide containing at least one amino acid differing from any of the known HCV types or subtypes amino acid sequences, or an analog thereof being substantially homologous and biologically equivalent.
- 42. A method for detecting antibodies to HCV present in a biological sample, comprising:
- (i) contacting the biological sample to be analysed for the presence of HCV with a peptide according to claim 41,
- (ii) detecting the immune complex formed between said antibodies and said peptide.
- 43. A method for HCV typing, comprising:
- (i) contacting the biological sample to be analysed for the presence of HCV with a peptide according to claim 41,
- (ii) detecting the immune complex formed between said antibodies and said peptide.
- 44. A. diagnostic kit for use in detecting the presence of HCV, said kit comprising at least one peptide according to claim 41, with said peptide being possibly bound to a solid support.
- 45. A diagnostic kit for HCV typing, said kit comprising at least one peptide according to any of claim 41, with said peptide being possibly bound to a solid support.

 SUBSTITUTE SHEET (RULE 26)

- 46. A diagnostic kit according to claims 44 or 45, wherein said peptides are selected from the following list:
- at least one NS4 peptide,
- at least one NS4 peptide and at least one Core peptide,
- at least one NS4 peptide and at least one Core peptide and at least one E1 peptide, or,
 - at least one NS4 peptide and at least one E1 peptide.
 - 47. A Diagnostic kit according to claims 44 to 46, said kit comprising a range of peptides which are attached to specific locations on a solid substrate.
- 48. A diagnostic kit according to claims 44 to 47, wherein said solid support is a membrane strip and said peptides are coupled to the membrane in the form of parallel lines.
 - 49. A pharmaceutical composition comprising at least one peptide according to claim 41 and suitable excipient, diluent or carrier.
- 15 50. A method of preventing HCV infection, comprising administering the pharmaceutical composition of claim 49 to a mammal in effective amount to stimulate the production of protective antibody or protective T-cell response.
 - 51. Use of a composition according to claim 49 in a method for preventing HCV infection as defined in claim 50.
- 52. A vaccine for immunizing a mammal against HCV infection, comprising at least one peptide according to claim 41, in a pharmaceutically acceptable carrier.
 - 53. A vaccine according to claim 52, comprising at least one peptide according to claim 41, with said peptide being unique for at least one of the subtypes or types as defined in claims 2 or 3.
- 25 54. An antibody raised upon immunization with at least one polypeptide or peptide SUBSTITUTE SHEET (RULE 26)

according to any of claims 22 to 26 or 41, with said antibody being specifically reactive with any of said polypeptides or peptides, and with said antibody being preferably a monoclonal antibody.

- 55. A method for detecting HCV antigens present in a biological sample, comprising:
- 5 (i) contacting said biological sample with an antibody according to claim 54,
 - (ii) detecting the immune complexes formed between said HCV antigens and said antibody.
 - 56. A method for HCV typing, comprising:
 - (i) contacting said biological sample with an antibody according to claim 54,
- detecting the immune complexes formed between said HCV antigens and said antibody.
 - 57. A diagnostic kit for use in detecting the presence of HCV, said kit comprising at least one antibody according to claim 54, with said antibody being possibly bound to a solid support.
- 15 58. A diagnostic kit for HCV typing, said kit comprising at least one antibody according to claim 54, with said antibody being possibly bound to a solid support.
 - 59. A diagnostic kit according to claims 57 to 58, said kit comprising a range of antibodies which are attached to specific locations on a solid substrate.
- 60. A pharmaceutical composition comprising at least one antibody according to claim 54 and a suitable excipient, diluent or carrier.
 - 61. A method of preventing or treating HCV infection, comprising administering the pharmaceutical composition of claim 62 to a mammal in effective amount.
 - 62. Use of a composition according to claim 60 in a method for preventing or treating HCV infection as defined in claim 61.

Figure 1		1		1	/74				
HCV-1	1 -	_	እ ር ር እ እጥ	רריים ז	ል ሮር ጥር ፣	מממממה	מהמממ		50 CACCAACCG
HCV-J									
HCG9	10					G	C		
BNL1									
BNL2									
CAM1078	10						C		
FR2									
FR2	TI								
HC-J6	2a	·	A- - -			G	C	A-A	·
HC-J8	2 h		2					2-2	A
S83									T
									T
NE92	20		A			G	C	A-A	<u>1</u>
FR4	21		A			G	CT-	A-A	T
BNL4	2 e		A			G	C	A-A:	r
BNL5	Zh		A			G	C	A-A	T
NET 1	2-		-7A C ITT			C	C	/ / / /	T
NZL1	3a		ACT-			G		A-A	T
HCV-TR	35		ACT-			G-C-	<u>C</u>	A-A	ACT
NE 48									T
NE 274									T
NE145									GT
NE125	3f		ATT-			G-C-	cc	A-A	ACC
						_	_		
Z4									
Z1									
GB358									
DK13	4 d					G	C		
GB809	4 e				T	G	c		
BNL7	4 k					G	c		
	_					_	_		
BE95	5a					G	C	A-A	
нк2	6.2		- ТСТ	4	C	6		A-A	
	Va		7301		· ·	O	Ū		
FR1	7a		-ACT-	A	C	G	C	A-A	T
**** <i>4</i>	_	,	T.O.T.	~	-	~	~	73 79	
VN4	Вa		-ACT-	A	0	G	·C	A-A	T
VN13	85		ACT-			G	C	A	
VN12	٥٠		- <u>አ</u> ርጥ	4		 C		Z -Z	A
AIAT	Эd		ACI	-A		- -	C	A A	Α
NE98	10a		-ACT-		A	G	c	A-A	N

F	igure 1 -con	tin	ued		2/74				
	HCV-1 HCV-J HC-G9 BNL1 BNL2 CAM1078 FR2	1b 1c 1d 1d	C1 C1	к-gs	T T NNNNNI N	1N	-CT	GTCAGATCG -C	C
	HC-J6 HC-J8 S83 NE92 FR4 BNL3 BNL5	2b 22d f e	C C		T		 -CT- -C	-C -C -C -C	C C C
	NZL1 HCV-TR NE48 NE274 NE145 NE125	3b 3c 3d 3e		A 5A	·T ·T		-C -C -C	-A -A -C -C	
	Z4 Z1 GB358 DK13 GB809 BNL7	4b 4c 4d 4e	C(C(C	CATT CAT -AT CAT	GA T T		-C -CT T	-C -C -C -C	C C C
	BE95 HK2								
	FR1	7a	5	CAT		·	-C	-C	
	VN4 VN13							-C	
	VN12								
	NE98	10a	CG		T	A	-C		

Figure 1 -	continued	3/74 .	
HCV-1 HCV-J	101 1a TTTACTTGTTGCCC	GCGCAGGGGCCCTAGATTGGGTGT	150 GCGCGCGACGAGA
HC-G9	16	CG CG	TG
BNL1	1d		
BNL2	1dC		CC
CAM1078	1e -CGC-A		<u>A</u> CC-C
FR2	1f		AGC-G
			G
HC-J6	2a -A		
HC-J8	2b	CG	AG
S83	2c -AC	·	
NE92	2d -A		
FR4	2f		C-AG
BNL3	2e		
BNL5	2h -A	CC-G	G
			_
NZL1	3a -AG	AC	C-T
HCV-TR	3b -ATGC	TAC	AGTAC-T
NE48	3c -AG	CT	TAC-T
NE274	3d -CAC	A	AGTTC-T
NE145	3e -A	AC	ATC-T
NE125	3f -AG-A	AC	AGT-C-T
Z4	4a	CG	TC
21			
GB358	4c	CG	TG
DK13			
GB809			
BNL7	4k	-	TC-G
BE95	C ~	~~	
DE 9 3	5a	GA	TC-G
HK2	6.5	22.5	•
nk2	6d	CC-G	
FR1	72		-
INI	74		
VN4	93	GC-C	-
VN13	8b		
11120			G
VN12	9a -C	AC-T	
	32 C -A	AC-1-÷	
NE98	10aGC-AA-	CCAG	TAGT-C-C

Figure 1 - con	inued 4/7	74	200
HCV-J 1 HC-G9 1 BNL1 1 BNL2 1 CAM1078 1	a AAGACTTCCGAGCGGTCGC. bA dG eG	TAG CGG TCG -GT-ACG	-A-AT -AT-T -AT-T
HC-J8 2 S83 2 NE92 2 FR4 2 BNL3 2	aC bAC cAAC dAC fTAC eTAC	-GGTAC -GATGG -GATGG -GATAG	-CCG -CCT -CCA -CCT
HCV-TR 3 NE48 3 NE274 3 NE145 3	aAAAAA	-GCAAACAG -GC-CGC-G CCAACC-G CCAC-G	C-T -G -G -AT
Z1 4 GB358 4 DK13 4 GB809 4	aGA	TCG TG TGG	
	aAC-		
	aCAC		
VN4 8 VN13 8	a	-GCAG -GCA-G	-CAAA -CAAG
VN12 9	aC	-GG-CAG-	-CAAA
NE98 10	a	G	-CACG

Figure 1 -	continued	5/74
	201	

	201
HCV-1 la HCV-J lb	250 GCTCGTCGGCCCGAGGGCAGGACCTGGGCTCAGCCCGGGTACCCTTGGC
	CCAATG
BNL1 1d	YYTTT
BNL2 1d	C-A-TTNNAC-TC
CAM1078 1e	-AGCAT
	CATA
	0 11 1 =A
HC-J6 2a	-AGCTACTAATGAA-AAAAC
HC-J8 2b 3	-AGCTACCA-TGAAAAT
\$83 2c 2	-AGCAACTA-TGAAGAA
NE92 2d 2	-AGCACTA-TGAA-AAAA
FR4 2f 2	-AGCGACTA-TGA-GTAA
BNL3 2e 2	-AGN-NGACTTGA-GTAATC
BNI-5 2h 2	- AGN-NGACI
DNII ZII Z	-AGCTACTAATGA-GTAA
NZL1 3a -	
	GAGACT
MCV-1K 3D -	CTCGCT
NE48 3c -	GTGGACTG
NE274 3d -	AAGCTT
NE145 3e -	AC-C-AGGAACTGTC
NE125 3f -	ACAAGCTCT
	GC-AAATG
Z1 4b -	GCTT
GB358 4c -	AAT-TATA
DK13 4d -	GC-AA-TTTTTT
GB809 4e -	GCATATGT
BNL7 4k -	GATATAAATA
BE95 5a -	GC-AACCTGA
· ·	
HK2 6a -	-GC-ACCAA
FR1 7a -	TAC-AGACAC-T-GGAC
, •	
VN4 8a A	TGC-AC-AAACC-TCC
	TGAC-AAACC-TAC
	10 AC AAAC C-1
VN12 9a -	TGC-A-AA-C-AC-ATC
72122 96 -	16 C A AA C -AC-A
NE98 10a -	-GCAAT
1.250 IUa -	G C M1

Figure 1- continued

			•	
HCV-1 HCV-J	251 la CCCTCTATGGCAATGAC lbC	GGCTGCGGGTGG	GCGGGATGGCTCC	300 TGTCTCCC
HC-G9	16	IAIG	A	A
BNL1	1c	1		C
BNL2	ld		N	C
FR2	1d	A		C
FK2	lfCTCT	A		C-T
HC-J6	2aACG	ACTC	A	C
HC-J8	2bGCAC	IC	T	C
S83	2cGG	CTC	AG	C
NE92	2dGCG	CTC	AG	C
FR4	2fGCGC	·CTC	A G	C -
BNL3	2eGGC			C-
BNL5	2hGGC			<u>-</u>
21.20	2.1. GC	0111	A	CT
NZL1	32		•	
HCV-TR	3aTC		<u>A</u> G	CA
	3bCGA	TT	AT-	C
NE48	3c			C
NE 274	3d -TT	T	A	C
NE145	3eTC		AGT - -	T
NE125	3fG	T	A	
Z4	4a		AG	 -т
Z1	4bTC	Т	AG	
GB358	4c -TTCT	T		
DK13	4dTC			X
GB809	4eTC		3 6	A
BNI.7	4 k - T . T . C . T		AG	CT
י דאדו	4 k -TTCT	T	T-	- - -C
מחסר				
BE95	5aTC-C	CT	AGG-	-CCT

HK2	6a -TTAC	T	-AT	C
FR1	7aTC	A		C
VN4	8a -TTA 8b -TTG	TT	-AC	C
VN13	8b -TTG	TTC	-AG	
VN12	9a T GC			C_
	<u> </u>			
NE98	102 7 6		n C	
NESO	10aAG		-AG	C- - G

Figure 1 - continued

				· ·
HCV-1 HCV-J	· 1a	301 CGTGGCTCTCGGCCTAGCTGG	GGCCCCACAGACCC	350 CCGGCGTAGGTCGCG
HC-G9 BNL1 BNL2	1c 1d	C	TT-T	GA
FR2	1f	CCT	AT	AAA
HC-J6 HC-J8	2a 2b	ATCTCTCT CGTCT	AT	AC AAA
S83 NE92	2c	CTCTCA AGCGTCA	C	AAA-
FR4 BNL3 BNL5	2e	G	A-AC	ACA
NZL1		CCTATC		
HCV-TR NE48		TT CT		
NE274		CATCT		
NE145	3e	CCAGT	AC	AC
NE125	3f	T	A-AT	AA
24	4 a	CATCT	A-ATT	GA
21	4b	CTCAGTCT	<u></u> ATT	<u>-</u> <u>C</u>
GB358	4 c	AGTCT	A-ATT	AC
DK13		GTCT		
GB809 BNL7		CT		GC
BE95	5a	AAT	TA	A-AA
HK2	ба	CACAT	TA	C-AC
FR1	7a	CGAT	AC	AC
VN4	8a	CCA-AT	A-AC	C
VN13		-NC		
VN12		CGGA	NAT	N-GC
NE98	10a	C		

Figure 1 - continued

351 400 la CAATTTGGGTAAGGTCATCGATACCCTTACGTGCGGCTTCGCCGACCTCA
1b TAA
1cT
1fTT
2aCGTT
2bTTT
2cc
2dCTTT
2fCT-S
2eN-NT
.3aAAAAA
3bcT
3cG
3dCCAAAT
3eTT
3fCTAT
4aC
4b TCA
4ccT
4dC
4eCCAA
5a TAT
6a GTTTT
7aC
8aCA
8bCC
9aCC

Figure 1 - continued

HCV-1 HCV-J HC-G9 FR2	1b 1c	401 TGGGGTACATACCGCTCGTCGGCGCCCCTCTTGGAGGCGCTGCCAGGGCC TTC-A-GA-GA-T TTA-GTAA	
HC-J6 HC-J8 S83 NE92 FR4 BNL3	2b 2c 2d 2f		
NZL1 HCV-TR NE48 NE274 NE145 NE125	3b 3c 3d 3e		
Z4 Z1 GB358 DK13 GB809 BE95	4b 4c 4d 4e	ACAG	
HK2		TCAGCAGTCAT	
FR1		CTGC-AA-GGGCTGGCT	
VN4 VN13	8a -	TCTGATGW-GTCGGN	
VN12	9a -	ACTGTCTGGCAA	

Figure 1 - continued

HCV-1 HCV-J HC-G9 BNL1 BNL2 FR2				-A-	- - -	-T-	-	GGTTCT AC		-G 			-T-	-C-	 	
HC-J6 HC-J8 S83 NE92 BNL3 FR4 BNL4 BNL5 BNL6		2b 2c 2d	C-	-A- -C- 	-C- 	-T- -G- 	-TA -GA -GA -G	AC C 		-G -G -G -G	GA GA GA	- A - - A - - A - - A - - A -	-T- -T- -T- -T-	-C-	 N	-G
NZL1 HCV-TR NE48 NE274 NE145 NE125	•	3b 3c 3d 3e	C- C- C-	-T- -A- -A-	 -c-	-T- -T- -G-	-GA -GA -GA	CC CA C A-CC C	T-0	GG -G -G	A- GA AA	 -T- -A-	 T-	TC- TC- T	 	
24 21 GB358 DK13 GB809 BNL7 BNL8 BNL9 BNL10 BNL11 BNL11		4abc4de4k4k4k4k4k4k4k4		 -A- 	 -C-	 -T- 	A	C-G- CCG- C-G- -C	- -	-G -G -G 	AA GA G-	-T- -C- -C- -C- -T- -T-	- T	-C- -C- T		
 BE 95		5 a	c-	-A-	-c-	-T-	-GA-	c	-T-	-G- - -	G - -	-A-				
нк2		6a	c-	-A-			-GA	CAA-	-C-	-G	GA	-c-	-T-			
FR1		7a			- 		-AT-	CAA-	-c	-G	G-	-c-	- - T-	-c-		-
VN4		Ва	T	-		-G-	AN-	NCA-	-C-	-Ģ	- - N	-A-	-T-	-C-		N
VN12		9a		NA-		-T-	A	CCA-	-C	-G	GA-	-A-	-			
NE98		10a									AA-	-T-	-T-	TC-		

11/74

Figure 1 -continued

		501					550
HCV-1	la	GAACCTTCCT	IGGTIGCIC	TTTCTCI	ATCTTCCTTCTG	GCCCTGCTCT	CTT
HCV-J	1b	TGC	<u> </u>		CT-A	TTG-	
HC-G9	1c	CC		T	T-GC	TTA-	-C-
BNL1					CT		
BNL2					CT-A		
FR2	1 f	N	J _	NN	CT	NT_7 _ G	
1112			•	1414		M1-W	
HC-J6	2a	T-AC		^T_	T-G		-C-
HC-J8					TT-GT		
S83	25	TT-CC	;_		TCT	11 - -G-	-A-
NE92							
					T-AT		
BNL3					TNGT		
FR4					T-G		
BNL4	2g	TG			T-GT	TG-	
BNL5					T-G		
BNL6					T-A		
DREU .	2.1	Ü	`	- 1	1-A	1-	-
NZL1	٦a	T-GC	·	~	T-	T	
HCV-TR					TCC-		
NE48							
					<u>T</u> -G <u>T</u> -		
NE274					T-GT-		
NE145					T-GT-		
NE125	3£	TT-GC	:	CT	T-	CI	-A-
24					T		
Z1					TT-		
GB358	4 c	TC			T-CT	-ATT	-G-
DK13	4 d	TC			CT	-A	-G-
GB809					CT		
BNL7					CT		
BNL8					CT		
BNL9					CI		
BNL10					Y		
BNL11	4 k	XCC			CT	-AT	-G-
BNL12	41	CC			A-C	-AT	-G-
BE95	5a	TT-AC			ATT	-TT	-G-
HK2	6a	TCC			T	-AA	-G-
		_					
FR1	7a	T	C	T	CT-A-	-AT-A	-G-
T757.4	O =	m ~	1757				_
VN4	ва	rC·	NN	N	NCT	-AT	-G-
TD11 0	^						_
VN12	Уa	T			WCT	-AT	-G-
NE98	100					mm m	70
いじつひ	TOG	T_M				-111	.H-

Figure 1 - continued

```
600
         1a GCTTGACTGTGCCCGCTTCGGCCTACCAAGTGCGCAACTCCACGGGGCTT
HCV-1
         1b -T----GA-C--A----C-T---G-G------GTGT-C---A-A
HCV-J
         1c --C---A--C--T------GT-GG------TT-----G-G
HC-G9
         1d -----G-T--AA-KA-C--TC--G-G------G-AT-C---G-G
BNL1
         1d -----G-T-AA--A-C--TC-TG-G-----G-AT-C---G-A
BNL2
         1f --C-C--A--C--A-C--T----TG-G----A--G-A-A--C-ATGGC
FR2
         2a --A-C--CACC--G-TC--C--TGC-G----AAG---AT--GTACCGGC
HC-J6
         2b --G-C--A----A-TG--T--AGTGG----CA-G--ATT-GTTCTAGC
HC-J8
         2c --A-CT----A-T---C---GTGG-G--CAAGG--A--GGC-ACTCC
S83
         2d -TA-C-----G-TC--C-G--TG--G--CAAG---A---GCA-CTC-
NE92
         2e -TG-C--C---T-TC--T-N-GTTG-G--CAAA--TA---GTCA-GCC
BNL3
         2f -TA-C--C----TG--T--ATA--G-TAAG---AA--GCCACT-C
FR4
         2q -TG-C--C---T-TC--T---GTG--G--TAAG---A---GTACCA-G
BNL4
         2h -TC-C------G-CAG--G--CAAG---A---GCCACTC-
BNL5
         2i --A-C--C----G-TC--T---GTG-----TGCG---CG--GT--TTC-
BNL6
         3a ----A-T-CAT--A--AG-CAGTCTAG-GTG---G--TA-GT-T--C--C
NZL1
         3b -----TGC----G--TAG-GTACACG---A-GT-T--C--A
HCV-TR
         3c ----GTCTGT--T--AG-A-GGCT-G-GTAC--G--TGTAT-C--C--C
NE 48
         3d ----GTCTGT--T---G-A-GGATTG--TAC--G--TGTGT-T--C--C
NE274
         3e ----CT-TGC--T--AGTC-GG-TGG-G--T-----G-AT-C--T--C
NE145
         3f ----GT-TCC----AG---GGCTAG-GTACA-G---A-GT-C--C--A
NE125
         4a --C-C----T--A--G----TG-G--CTAC--G--TG-TT----CA-C
24
         4b --C---AACA--A--A--T---GTG--CTAC--G--TG-TT----CG-C
21
         4 c -- C------T---A-C------GT-A-CTAT-----TG--T-----CA-C
GB358
         4d --C----T----TG-C
DK13
         4e --C-C----T-----G----G-GTTA-CTAT-----TG-TT----CG--
GB809
         BNL7
         BNL8
         BNL9
         BNL10
         BNL11
         41 --C------C--G--C----TC-G--TTAT--G--TGT-T----CA--
BNL12
         5a -TC----C--T--G--C--T--AGTT-CCTAC--A--TG--T-T---A--
BE 95
         6a --C-C--AAC---A------TCTTACCTACG------A
HK2
          7a --C-C---ACA--A--C--A--AATT-----CAAG---G--T-T---A-C
FR1
          8a --C-T--AACA--A--C--C--GGCG--TTATAC----AAGT-T--C--G
VN4
          9a --C-C--CAC---T--C--C--ACTAA-CTATGCT---AAGT-T----G
VN12
         10a ----CT-ACA---A-AG-C-GGCTGG-GTAC--T--TG--T-C--A--C
NE98
```

Figure 1 - continued

HCV-1 HCV-J HC-G9 BNL1 BNL2 FR2	601 1a TACCACGTCACCAATGATTGCCCTAACTCGAGTATTGTGTACGAGGCGGC 1bTGCCT-CATA 1cTCTGTCCGAA 1dTTCTT-CCCA-CTATA 1dTTCCTT-CCCA-CTAT-AG 1fTTCTT-CGGCCCATAAA
HC-J6 HC-J8 S83 NE92 BNL3 FR4 BNL4 BNL5 BNL6	2aATGGCCA-CTGATCACC-GGC-ACTCCA 2bTCTT-AAACCCACC-GGCCTCA- 2cATGCCGCT-CT-CT-GGCCTT-A 2dATGACAGAGTCCC-GGCCTCAG 2eTATG-CACT-CAACCCA-GGC-ATTN 2fATG-CGTCTG-CTGACCCC-GGCCTCAG 2gATG-CACTT-CAACCCA-C-GGC-AAT-CA 2hTATGGT-AAGCCC-GGCCTTAA 2iATGGT-GAGCCC-GGCCTC-A
NZL1 HCV-TR NE48 NE274 NE145 NE125	3a GT-C-TC-CTT-CTAGCTC-A 3b TGTGC-TC-CTTGGCC-A 3c ATACC-TT-GAGCCATC-A 3d GTGCCTGGCCTCC- 3e ATGCCT-AAGCCAA-TA 3f ATAC-TCCTAGCCCTT-
24 21 GB358 DK13 GB809 BNL7 BNL8 BNL9 BNL10 BNL11	4a TA
BE95	5aTTT
HK2	6aTCACC
FR1	7aTC-TCT-GAACCCT-TTA
VN4	8aTC
VN12	9aTTC-ACTAGCC
NE98	10aATGATCCAGGGTCTC-G

Figure 1 - continued

```
651
                                                          700
           1a CGATGCCATCCTGCACACTCCGGGGTGCGTCCCTTGCGTTCGTGAGGGCA
HCV-1
HCV-J
           1b G--CATG---A------C--C-------G--C-----C--G---A-T-
           1c GA-CCTG---A----TCTG--C----T--G--C-A---A--C----
HC-G9
           1d --G-ATG---A----TAC--A-----G--C-----G----AT-
BNL1
BNL2
           1d T-G-ATG---T----G-C--A----T--G--C-----G---AA--
           1f G--CAT-----T-----G--T----N--G--C--A-A--G--A----
FR2
HC-J6
           2b T--C--AG-T--C--TCT---T--A----A--T-AGAA---TAATG
HC-J8
S83
           2c A-GA--AG-G--T--T-----T-A-----T-AG---ACC-C--
NE92
           2d G----TG-T--T---GTC--C----T----T-AGGAGA-----
           2e G--C--GG-G--T--TGT---T-A--T----C---AGAA-AGCTC-G
BNL3
           2f G--C--GG-G--C--TGT---T--A--T----C--T-AGA-GTCA--T-
FR4
           2a G-GC--GG-G--T--TGT---T--A--T----G--T-AGTTGC----
BNL4
           2h G----TG-G--T---GTC--T--A--T--T--A--T-AGA-GC-CCAA-
BNL5
          2i G--G---G---T---GTC--T--A--T--T--C--T-AGT-GA---A--
BNL6
NZL1
          3a T---T--T---A--C--C--T--A----T--C-AG--C----
HCV-TR
          3b A----TG---T----TTA--C--A----G--C----CACAACC----
NE 48
          3c -C---T----T----TTG--C--T----A--C----C-AAA-CAAT-
NE274
          3d T--A-T----TTG--A--T--T-G--C-----AATCA----
NE145
          3e A----TG------TG--T--T-----T--C-----G-AGA-C----
          3f TA---T-----TG--C--C--T--G--C--AC---C--T-
NE125
24
          4a -C--CA-----A---TTG------A--C--T--GATGACT--G-
z1
          4b GC-CCA----A----TTG--A----T----C--T--G--GAC--AG-
GB358
          4c GC-CCA----A---CTC-A----TT-A--C----GA-G-TT--G-
DK13
          4d TT-CCA----T-A---CTC----A----T------GA-G--A--G-
          4e -A--CA----T-A---CTC--A-----A--C--T--GAAGACC--G-
GB809
          4k -C--CA----T-----CTC--A--T-----G--C------GA-A-----G-
BNL7
          4k -C-CCA----T-----CT---A--T-----G--C-----GA-AACT--G-
BNL8
BNL9
          4k -C--CA----T----TCTC--A--T-----G--C------GA-A-T---G-
BNL10
          4k -C--CA----T-AGCACT---A--T----G--C-----GA-A-T---G-
BNL11
          4k -C--CA----T-----CT---A-A-----G--C-----GAAA-----A-
BNL12
          41 -C--CA----T-A---CTA--A----T--A--C--T--GAAGACT--G-
BE95
          5a TA-CCTG----A---G-A--T--T----G----T--CATGACA--T-
HK2
          6a T-C-ATG---T---TTTG--T--A---T-GA-G-TC-ATG
FR1
          7a GACCATG--A----TCT---A--T--T---A--TA-CAAG-C---G-
          8a GACACTG--TT-----TTG--T----T-A----T-GAAGRT-RA--
VN4
VN12
          NE98
         10a G---ATT-----C---TTA--T--C--T-----C----A--CTCT----
```

Figure 1 continued

```
750
HCV-1
           1a ACGCCTCGAGGTGTTGGGTGGCGATGACCCCTACGGTGGCCACCAGGGAT
HCV-J
           1b -TTT---CC-T--C---A---C-C--T--C---C-C--GG-----A-C
HC-G9
           BNL1
           1d --CATCTCC-C--C--A----C-C---C-T--GGT--AAA-Y
BNL2
           1d --T-T--TC-T--C---A--C-RC-C-----C-T--GGT--AA--C
FR2
           1f -TAT---CC-T--C----AC--C-C----C--C-C-AG-GC--ATC
HC-J6
          2a -TA-A--TC----C--A-AC--G-CT-A--G-AT-----GTGCA-C-G
          2b G-A---T-CAT--C--A-ACAAG-A--A--C-AC----TGTG-AAC-C
HC-J8
S83
          2c ---T---TC-A-------C--G-TG----C-ATC-C----TA--TC-A
NE 92
          2d --ATA--CC-C----A-AC--G-TT-G--C-ATA-A--TGTG--CC-A
          2e GTCGG-TCCAC----A-CC----CT-G--C-ACA-A---GTG--CA-A
BNL3
FR4
          2f -TAGGA-CTTC-----ACA---G-CT-G--C-AC----TGTG--CCGA
BNL4
          2g -TAAG--CC----C--A-AC--G-C--T--C-AC----TGTG-ACC-G
BNL5
          2h -TCAG--TC-C--C--A-AC-TG----A-C-AT-----GTG--CC-A
BNL6
          2i --A----CC-C--C--A-AC--G-C-----ACA-C--TGTG--CC-A
NZL1
          3a - TA-A--T-C---C---ACCC-AG----A----A----AGT----T-C
HCV-TR
          3b -- CAA--ATCA--C---ACAA--G-CT-AA-G------GTT---ACC
NE48
          3c --A--A---C---C---A-AC--G----T--G--A-----GGT---TC-C
NE274
          3d --T----CAA--C--A-TC--G--G-A--A--A-----GGTT-A-T-C
          3e --A-A---GA---C---ACCC--GC---A--A-----AGT---AT-C
NE145
NE125
          3f -- CAG--A----C--AC-C-AG-A--A--G--A----TGT--AAC--
          4a --A-A---C-T--C---AC-C--G----G----A----TGT-GCAC-C
Z4
21
          4b -TA-T--TC-C--C-----C-CT-----C--T-----G-GCCCT--
          4c -TCAG--AC-C--C-----CC-C--T--C--C-----GG-GCCTT-C
GB358
          4d --AAG--T-CA--C----T-TC-C----C--C--TG-GCAAC--
DK13
GB809
          BNL7
          4k -TCAG--AC-T--C---A--CC-T-----C--C--AG-GCCAT-C
BNL8
          4k -TCAG--AC-T--C-----CC-T--T-----C--C--AG-GCCAT-C
          4k -TCAG----T-C-----CC-T-----CA-C--AG-GCCAT-C
BNL9
BNL10
          4k -- CAG--AC-C--C------CC-T------C--G--GCCAT-C
BNL11
          4k -TCAT--AC-C--C------CC-T-----C--C-AG-GCCAT-C
BNL12
          41 --A-T---C-C--C------CT-A--A-------G-GCCCATA
BE95
          5a -T-TGAGT--A--C-----CCAA--T-----AC--T-AG--CC-AGC
HK2
          6a -TCGG--C-CC-----CAT--TG-----C--CC-----TACCAA--
FR1
          7a -T-AG--AC-A-----C-CC-TG-CT----C-CT-A---GT-CCCA-C
VN4
          8a -TCAA--CC----CA-GCCT----G--CC---AGTGCC-A-C
VN12
          9a --CTGA-C-A-----C--T--GCCT----G-AT----GGTGCA-A--
NE98
         10a -TA-A--A--C--A-CC-TG---G---Y--C--GTG-A-TCG
```

Figure 1 - continued

```
751
                                                             800
HCV-1
           1a GGCAAACTCCCCGCGACGCAGCTTCGACGTCACATCGATCTGCTTGTCGG
           1b A---GCA----A-C---ACAA-A-----C---G-----T----C--T--
HCV-J
           1c TCGCGCG-----TC-GTG--G----GTG----CTC-A-----
HC-G9
           1d -CT-GTG----A-TR--GCAA-C-----G----CT-----T--
BNL1
BNL2
           1d -CT--TG----TA-TG--GCAA-C----CT----G--T--
           1f -CG--CGCT---ATCGATG--G-G--G----G---C--C--C--C--
FR2
HC-J6
           2a CC-GGCGC-T-A--CA-GGCT-A--GACG----T--CA--G---GAT
HC-J8
           2b C--GGTGCG-T-A-TCGTAGC--G---ACA---G---CA--A-C--AAT
S83
           2c CCTGGCGCT-T-A-T-A-GGC--G---GCA----A-CA-C--GAT
NE92
           2d CCTGGTGCG-TTA-C-A-GGC--G--GACG--T--T--ACCA-CA-T-C
           2e CCTGGTGCT-T-A-C-A-GGA--G--GGCA-G---T---GCCG-C--GAT
BNL3
           2f CCTGGTGCT-T-A-T-GAGGT--G--GGC----T--ACCA-C--GAT
FR4
           2q CC-GGCGC-T-A-T-G-GGCT-G--GACG----T-CACCA-C--GAT
BNL4
           2h CCTGGCGCG-T-A-C-G-GGTT-G--GACG----T-CACCA-C--T-C
BNL5
BNL6
           2i CCTGGCGCG-TTA-C-A-GGC--G--GACA--T--T--CA-CA-----C
           3a -T-GG-GCAA-TA-TG-TTC-A-A--CA----TG-G--C--AT-A--A--
NZL1
           3b CTTGGCG-GA--A-CG--TC-A-C--ACC--TG-G---A---G--A--
HCV-TR
NE48
           NE274
           3d -CTGGCGCGA--A-TG-ATC-A-C--CA----TG-G-----G--G--
NE145
           3e -CTGGTGCAA-GA--G-TTCCG-A--CG-A---G-G---T----A----
NE125
           3f CCTGGCGCAGT-A-CG-ATCAA-C--CA-G--TG-G---T-A-G--G--
24
           4a CCGGGCGCT--GCTTGA-TC-T-C--G--A--TG-G--CT-AA-G--A--
z_1
           4b CC---CGCA--GTTAGA-TCCA-G--CA-G--TG-A--C--A-G--G--
GB358
           4c AT-GGCGCT--GCTTGAATCC--C--GA----TG-G-----A-G--A--
DK13
           4d CTG--TGCT--GCTTGA-TCTT-GA-----G-G-----A-G--G--
GB809
           4e -T-GGTGCT--GCTCGA--CCT-G--G--C--TG-G--C--A-G--A--
BNL7
           4k AT-GGCGCG--ACTTGA-TCT--A-GA----TG-G--CT--A-G--G-
BNL8
           4k AT-GGCGCA--GCTTGA-TCT--G--GA----TG-G-----A-G--G-
BNL9
           4k AT-GGCGCA--GCTTGA-TCCT-G--GA----TG-G-----A-G--G--
           4 k AC-GCGGCG--GCTTGA-TCC--G--GA----TG-G----A-G--G--
BNL10
BNL11
           4k AT-GGCGCG--ACTTGA-TCT--A--GA----TG-G---G--A-G--G--
           41 CTTTCGGCT--ACTT-T-TCCG-A--G--G--TG-G-----A-G--G--
BNL12
BE95
           5a CT-GG-GCAGT-A--G-T-CT----GA-AGC-G-T--CTAC--A-CG--
HK2
           6a -CTTCCACG----A---GGAT-C--CA-G--TG-G----T----CG--
FR1
           7a TCATC-G-G--AATCCACGG-T----C--A---G-A--C--C--C--T--
VN4
           8a -CGTCTACG--A-TC--CGG-T-C--CAAA--TG-G--CA-CA-G--G--
VN12
           9a -CGTCGG-GT--ATC-G-GGTG-C--CGAG---G-G--C--CT-G--G--
NE98
         10a CC-TGCGC-G--A-CG-CTCT--C--CACG---G-G---A-A-G--G-
```

```
Figure 1 - continued
```

HCV-1 HCV-J	801 1a GAGCGCCACCCTCTGTTCGGCCCTCTACGTGGGGGACCTATGCGGGTCTG 1b -GCGTG-TCTA-GTTCAC-	
HC-G9	1c -GCTG-GTTA-GTA	
BNL1	1d -G-NNGTCTA-GRT	
BNL2	1d CAG-GT-TCCTA-GCAC-	
FR2	1f -GCAGTGTCAGA-TTTGGC-	
	• 1	
HC-J6	2a -TCGCCTT	
HC-J8	2b -GCATGGCCT-GTATG-GG-C-	
S83	2c -TCTTGGTTTG-GTCG-GC	
NE92	2d ATCTGT-TCTGA-A-A-AAGTCG-G-	
BNL3	2e -TC	
FR4	2f -TC	
BNL4	2g -GTTGTAA-CG-GTCG-G-	
BNL5	2h -TCTTGCATT-GTCT-CCG-A-	
BNL6	2i -TCGTC-TT-GT	
	21 10 G1 C11-G1	
NZLI	3a CGCGGA-GCTGTTA-GTG	
HCV-TR	3b CCCDCDCDCDCDCDCDCDCDCDCDCDCDCDCDCDCDCD	
NE48	3b CGCACGACAAGGGCGCT-TG	
NE274	3c T-CGTAT-GATCTG-A-	
NE145	3d AGCTTGT-GCCGGTTCTA-GTAG-C-	
	3e CTTGCCGTCTG-C-	
NE125	3f TGCAG-GA-ATT-GG	
7. 4		
Z4	4a CGCGTT-GTTTCAGG	
21	4b TGCGTTA-GCTA-TA-TGTAGGC-	
GB358	4c TGCTTGCGCCTTA-CAGTGGC-	
DK13	4d CGTCCA-CAG-GTGG	
GB809	4e TGCTG-GCCCCGTGGCT	
BNL7	4k -GCTG-TATA-CTT-RTYGGCT	
BNL8	4k -GCTTG-TC-ATA-CTT-GTCGGCT	
BNL9	4k -GCGTGATA-CTT-GTCGC	
BNL10	4k AGCTTG-TATA-CYT-GTCGCCT	
BNL11	4k -GCTTG-TAT-GCCT	
BNL12	41 TGCATA-CGTT-ACGG	
•	1 A C = ===GG==	
BE95	5a AG-GTGCCGT-AAAGCGTG-AC	
	ou no o le Grand-AGCG1G-AC	
HK2	6a CGCAGTGG-TCATGA-CGTCC	
FR1	7a -GCAGG-AT-TA-GA-CA-CC-TTAGCA	
VN4		
* 44 2	8a CGCTG-GTATA-GTGGCC	
VN12		
A1417	9a TGCTTG-GTCTA-GCTTGGGC	
NE98	10a DCCC	
HESO	10a RGCGACATAAATTAG-GC	

18/74

Figure 1 - continued

```
851
                                                 900
HCV-1
         1a TCTTTCTTGTCGGCCAACTGTTCACCTTCTCTCCCAGGCGCCACTGGACG
HCV-J
         1b -T----C---TC---G-------A--TC-C--GT-TGA----
         HC-G9
         1d ----C-CT----G-A----T--A---C-CATG---CAT--A
BNLl
         1d ----C-CTTGT--CAT--A
BNL2
         lf ----C--C--T--G---T------A-GT--C---G--T-----
FR2
         2a -GA-G----CA-C---GA-----TTG----G--ACA--A-----TTT
HC-J6
         2b -GA-GA--C-ATCG--GGCT----TGG-A--A-ACAA-----AACTTC
HC-J8
         2c -GA-G--G-C--CT--GG-CG--GT-G-G--G--ACAA-A---TAC-TTT
S83
NE 92
         2d -GA-GT-G-CTTCT---G-C---T-A---G--CA--AT--TAA-TTT
BNL3
         2e -GA-GA-A-CT-CA--GGCT----T-G-GG-A--G-A----T-ACTTC
FR4
         2f -GA-GA-A-CA-CG---G-TGC-GT-G---A--GCAATA---TACTTTT
BNL4
         2g -GA-GA-A-CT-CT--GG-TG---TTG----G--GCAA-AT---AACTTT
         2h -GA-GT-G---TCT---T-T---TGA----C--TCA--A----ATCTTC
BNL5
NZL1
         3a -----C--G--A---GCC-----G---AGA--TC-A-----TCAA---
HCV-TR
         3b -G------AC---C
NE48
         3c -T--C--C--A--A---GCA-----A---AGA---C-A----CA---A
NE274
         3d ----CT-G--G--A--GGCT------AGA--TC-T-AG---AAC---
         3e ----C-----G--G-GGCC--T--A---AGG--TC-T--T--TAC---T
NE145
         3f -T--C----G-----GC-----T---AGAG-TC---AA--T-AT--C
NE125
Z4
         4b ----C--A--G-----GA----CGA--GC-C--G-------C
21
GB358
         4c -A---T-G--T--T--GA-----T-T---CAG--GC------T
DK13
         GB809
         4 k -G--C--A----T--GA-----T-T---CGA--A-----T
BNL7
BNL8
         4k -G--CT-G--T--T--GA----TT-T---CGA--AC-A-----T
         BNL9
BNL10
         4 k -G--CT-G--T--T--GA-----T-T--YCAG--TC------T
BNL11
         4k -G--C--G--T--T--GA-----T
BNL12
        BE95
        5a -A--CT-G--A-----A------ATAGG--TC-C-AG---GCT---
        6a ----T-G-CG-A----A----TCAG---C-C-T-T----T
HK2
FR1
        7a -AA-CT-G--A--G--G--T--T--T--AGG--T-A-TA---TCA-GTT
VN4
        8a -T--C--C--T--A--G--C----GC--AGG--TC--ATG--TCA-GTT
VN12
        9a -----C--T--G--GT------G---AGA------ATGT-TGA--TC
NE98
       10a -A-----Y-G-GGG----T-A-GGAGA-ATC-C-AG--T-----T
```

Figure 1 - continued

```
901
                                                 950
HCV-1
         1a ACGCAAGGTTGCAATTGCTCTATCTATCCCGGCCATATAACGGGTCACCG
HCV-J
         1c -----G-G--A----T--
HC-G9
BNL1
         1d -----G-AG------A---
         ld --A--G-AG-----A---
BNL2
FR2
         HC-J6
         HC-J8
         2b --C---AG-----C-T--C-----C-AA--T--C--C--C--T--
S83
         2c GTC--G-AA-----C--T--C--A--C--GC--T----A----
NE 92
         2d GTC--G-AC-----C--T--C--A--C--A-----C--C--T--A--T--
BNL3
         FR4
         BNL4
         2g T-C--G-A-----T--C---
BNL5
         2h GTC--G-A----C---G--A
NZL1
         3a GTC--GACC--T--C----GC-G--C--A-----C-TT-A--A--T--
HCV-TR
         3b GT---GACG-----C----G-A--C--A------G-TT-A--A--T--
NE48
         3c GTT--GCA-----C----AC-G--C--A--T---G-TT-A----T--
NE274
         3d GT---GACC-----AC-G--C--T--T--C---T-A--A--A-
NE145
         NE125
         3f GTC--GTTG------AC-A--C-A--C--A--A--C--T-A-A--T-A
Z4
         4a ----G-AG-----T--C----CA-T-----C--C--C--A-
21
         4b --C--G-A-----C----C-----T--T--CG-CT----C--A-
GB358
         4c ----G-AC-----T--C----CG-G--G--CG-T-----C--A-
DK13
         4d --C----AC------T--C-----CA-A--A--A---C--A--A--A--A-
GB809
         4e --C--G-AC--T----T--C----CG-A--G----T----C--T--
BNL7
        4k --T----A-----T--C---
BNL8
        4k G-C--G-A-----T-----
BNL9
        4k -- C---- A----- C----
BNL10
        4k --C--G-A-----T--C---
BNL11
        4k --C--G-AA-----T--C---
BNL12
        41 GTC----AC-----C--T--C---
BE95
        5a GT---GAAC----C--T--C--T--CAGT------G-T--C--C----
HK2
        6a GT----AC-----C----A-A-----CG-C--C--C--A-
FR1
        7a --C--G-A---T--C-----NA-CN-T-----CG-C----A---A-
        8a GTC--G-AG--T--C--T--C-----CA-A--G-----C--T--A----
VN4
VN12
        9a G-C--G-AC----C--T--C-----G-A-----C--C--T--G----
NE98
       10a GTC--G-AC----C--T--C---
```

Figure 1 -continued

HCV-1 HCV-J HC-G9 FR2	951 957 la CATGGCA lbT lc AT lf NNNNNN
HC-J6 HC-J8 S83 NE92 BNL3 FR4	2aG 2bT 2cG 2d GG 2eG 2f ANN
NZL1 HCV-TR NE48 NE274 NE145 NE125	3a AT 3b TG 3c GT 3d GT 3eT
Z4 Z1 GB358 DK13 GB809	4a GG 4b GC 4c G 4d AT 4e GT
BE95	5a G
нк2	6a GT
FR1	7a G
VN4	8a A
VN12	9a GG

Figure 2

50 GVYLLPRRGPRLGVRATR	1 MSTNPKPQKKNKRNTNRRPQDVKFPGGGQIVGG	1a 1b	HCV1 HCV-J
X	R-TXXXXX	1d 1d 1e	BNL1 BNL2 CAM1078
A-	R-T	1f	FR2
	R-T	2a 2b 2c 2d 2e	HCJ6 HCJ8 CH610 NE92 BNL3
P-	R-T	2f	FR4
VV-	LRQTLN	3b	HCVTR
	R-TM R-TM	4 d 4 e 4 e 4 k	DK13 CAM600 GB809 BNL7
M	R-T	5a	BE95
	LR-TT	ба	HK2
	LR-TM	7a	FR1
	LR-TI	8a 8b	VN4 VN13
	LR-TM	9a	VN12
VV-	LR-TX	10a	NE98

Figure 2 -	- continu	ned 100
HCV1	la '	51 KTSERSQPRGRRQPIPKARRPEGRTWAQPGYPWPLYGNEGCGWAGWLLSP
HCV-J	1b	X-XSXXXX
BNL1	1d	DQSD-XXH
BNL2	1d	
CAM1078 FR2	le 1f	
	_	LDST-KS-GKLL
HCJ6	2a	
HCJ8	2b	
CH610	2c	
NE92	2d	
BNL3	2e .	L
FR4	2f	
HCVTR	3b	KQ-HLSRSKKL
	A -3	QLS
DK13	4d 4e	TS
CAM600		
GB809	4e	X
BNL7	4 k	
BE95	5a	AL
нк2	6a	Q-QH
FR1	7a	V-Q-TS-G
	0 -	V-HQT
VN4	8a	V-HQT
VN13	8 <i>p</i>	
VN12	9a	AV-QNQ
NE98	10a	SRTS

	_			23/74
Figure	2	_	continued	23/ 14

-		
HCV1 HCV-J BNL1 BNL2 FR2	la lb ld ld lf	101 150 RGSRPSWGPTDPRRRSRNLGKVIDTLTCGFADLMGYIPLVGAPLGGAARANNS-T
HC-J6 HC-J8 CH610 NE92 BNL3 FR4	2a 2b 2c 2d 2e 2f	VV
HCV-TR	3ъ	NF
GB116 DK13 CAM600 GB809 G22 GB549 GB438 BNL7	4c 4d 4e 4e 4f 4g 4h 4k	VV -XXNX
BE95	5a	NNK
HK2	6a	HNVV
FR1	7a	NNVL-GVL-GV-A-
VN4 VN13	8a 8b	NNXXIE
VN12	. 9a	D-X-NXV-AE
NE98	10a	N

Figure 2 - continued

HCV1 HCV-J	1a 1b	151 200 LAHGVRVLEDGVNYATGNLPGCSFSIFLLALLSCLTVPASAYOVRNSTGL
BNL1 BNL2 FR2	1d 1d 1f	XT-HEAS-V TT-HEAS-V -XXGXXXXXXXXXTE-HST-DG
HC-J6 HC-J8 CH610 NE92 BNL3 FR4 BNL4 BNL5 BNL5	2a 2b 2c 2d 2e 2f 2g 2h 21	F
HCV-TR	3b	A-GFFCGLEYT-TS
GB116 DK13 CAM600 GB809 G22 GB549 GB438 BNL7 BNL8 BNL9 BNL9 BNL9 BNL10 BNL11	4c 4d 4e 4f 4g 4h 4k 4k 4k 4k	-EAVI
BE95	5a	IVPYAS-I
HK2	6a	AII
FR1	7a	AIIK-AS-I
VN4	8a	XXIXX-X-X-X-X-XXTAHYT-KS
VN12	9a	-XAIIXTLNYA-KS
NE98	10a	I-FFLT-TAGLEYAS

Figure 2 - continued

HCV-1	1a	201 YEATHDCDMCCTABADD TAXIDD COMPANY 250
HCV-J	1b	YHVTNDCPNSSIVYEAADAILHTPGCVPCVREGNASRCWVAMTPTVATRD
BNL1	1d	SLL-A-N
BNL2	1d	SI-MDGM-M-YD-HLM-LL-VKX
FR2	1 f	LSI-MSGMAN-SMXLL-VK-
		S-GK-IXIPLL-A-I
HC-J6	2a	-MT-DTWOLOA-VVEKVTIPVS-NVOO
HC-J8	2b	-YAS-NTWQLTV-LENDNGTLHIQV-NVKH
CH610	2c	-MEQIPVS-NI-O
NE92	2d	-M
BNL3	2e	-MaS-NWOLKVEEK1IPVS-NI-VSQ
FR4	2f	-MAS-NWQLXVVENSSGRFHIPIS-NI-VSK
BNL4	2g	-MAA-DWQLRVVE-SRTFT-VS-NVSR
BNL5	29 2h	-MAS-NIWOMOG-VVELQKIPVNVNQ
BNL6		-MSWOLKVVF-HO-OTDVN- 1/60
DIATO	2i	-MSWQLEE-VVEWKD-TIPVNI-VSQ
HCVTR	3b	-VLS-GE-VLTTQ-STTVSTV-T
	-,	L-VL
GB116	4 c	IDYHLLVQLAPY
DK13	4 d	TDYHLK-TSLAQH
CAM600	4e	IATENHLTQLSPY
GB809	4e	I
G22	4 f	LFVHHLTQLL-APY
GB549	4 g	DHH-M-LTTPLAPY
GB438	4h	
BNL7	4 k	-VTPU T
BNL8	4 k	-YDHHLQLAPY
BNL9	4 k	
BNL9	4 k	I
BNL10	4 k	DHHALVQLAPY
BNL11	41	FDHHLKHLAPY
GB724		SDHHLKTTLAPI
GB 124	4×	IVTDHHLTVTPVAVS
BE95	5a	DNLAMTVQILSAPS
		ZNZ AQIDSAPS
HK2	6a	LLDAMLVDDR-TH-VL-IPN
FR1	7a	The second of th
	7 u	LS-NFETMLIKAELPVSL-VPN
VN4	8a	LETLL
10110	•	
VN12	9a	LNGMLKTLTKLSASL-VQN
NE98	10a	-MS-GG-ILSTIPVSXVKS

Figure 2 - continued

	251 300
HCV-1 la	GKLPATOLRRHIDLLVGSATLCSALYVGDLCGSVFLVGQLFTFSPRRHWT
HCV-J 1b	SSI-T-TIYE-
BNL1 1d	ASV-TXAIVXX-FMXAAM-H-
BNL2 1d	DNV-TDAIVT-AFRMLYH-
FR2 1f	ANA-TDEVVA-VFM-IGTS
HC-J6 2a	PGAI.TOGTMV-MG-M-AA-M-IVQHF
HC-J8 2b	RCALTRST-V-MI-MAAVA-MILS-A-MVONF
CH610 2c	PGTLTKGA-V-VI-MVALMIAA-AVIAQTF
NE92 2d	PGAI.TKGTTIAFIA-M-AS-V-IIOH-KF
BNL3 2e	PGALTKGARAV-MVA-MIAA-A-IVA-KYF
FR4 2f	PGALTRGATI-MIA-MIAA-VAVVQY-TF
BNL4 2q	PGALTRGTTI-MVIVA-MIAA-VVIVQH-NF
BNL5 2h	PGALTRGTTI-AVFA-MS-F-MIQH-IF
BNL6 2i	PGAXTKGTII-AF
22	
HCVTR 3b	LGVTTASI-T-V-MARQAF-AAAT-
GB116 4c	VGA-LESS-VMAVIGM-S-Q
DK13 - 4d	INA-LESVMGIVGQ
CAM600 4e	AGA-1.F.PVMAMIGLMQ
GB809 4e	VGA-LEPVMAVGLMQ
G22 4f	LGA-LESMV-M-TGI-A-MR-L
GB549 4g	VGA-LESMV-MAVIGMR
GB438 4h	LGA-L-SV-Q-V-M-AI-H-GA-MVS-Q
BNL7 4k	IGA-LESS-VMAVIX-XGLM-S-R
BNL8 4k	IGA-LESS-VMAVIGLM-S-R
BNL9 4k	IGA-LESS-VMAVIGAM-S-R
BNL9 4k	TAA-LESS-VMAVI-XGLM-SXQ
BNL10 4k	IGA-LESS-V-VMAVIGLM-S-R
BNL11 41	LSA-LMSVV-MASGAMQ
GB724 4x	VDA-LESFVMAVGAMQ
BE95 5a	LGAVTAPAV-Y-A-G-AAALMYRQ-A-
нк2 ба	ASTGFVA-A-VVSILAQ
FR1 7a	SSV-IHGFVA-AFM-IIIR-KY-QV
	TD M ON
VN4 8a	AST-V-GF-K-V-IMA-AFMGLLRM-QV
	CI
VN12 9a	ASVSIRGV-E-VA-AFMGLRMYEI
	PCAATAST-V-MM-XAALXG-SWRH-Q
NE98 10a	PUAATAST-V-MM-XAALI-ALI-AG-SWIMI Q
	·

Figure 2 - continued

Figure 2 -	COHLETIN	
HCV-1 HCV-J BNL1 BNL2 FR2	la lb ld ld lf	301 319 TQGCNCSIYPGHITGHRMA V-DVSE V-DSXXX
HC-J6 HC-J8 CH610 NE92 BNL3 FR4 BNL4 BNL5	2a 2b 2c 2d 2e 2f 2g 2h	V-D
HCVTR	3b	V-TVS
GB116 DK13 CAM600 GB809 G22 GB549 GB438 BNL7 BNL8 BNL9 BNL9 BNL9 BNL10 BNL11 GB724	4c 4d 4e 4f 4g 4h 4k 4k 4k 4k 4k	DAVDTDTD
BE95	5a	V-NSV
HK2	6a	V-DTV
FR1	7a	DXNXV
VN4	· 8a	V-ET
VN12	9 a	A-DA
NE 98	10a	A-D

Figure 3

SEQ. ID NO. 1 (BNL1, 1d)
ATGAGCACGAATCCTAAACCTCAAAGAAAAACCAAACGTAACACCAACCGCCGCCCTCAKGGSGTN
NNNNNNCCGGGTGGCGGTCAGATCGTTGGTGGAGTTTACCTGTTGCCGCGCAGGGGCCCCAGGNNG
GGTGTGCGCGCGACTAGGAAGACTTCCGAGCGGTCACAACCTCGTGGCAGGCGACAGCCTATCCCC
AAGGCTCGYCGGYCCGAGGGCAGGTCCTGGGCTCAGCCCGGGTATCCTTGGCCCCTCTATGGCAAT
GAGGGCTGCGGGGTGGGCGGGNTGGCTCCTGTCCCCCCGCGGCTCTCGGCCCAATTGGGGCCCC

SEQ ID NO. 3 (BNL1, 1d)
GACGGCGTGAACTATGCAACAGGGAACTTGCCCGGTTGCTCTTTCTCTATCTTCCTCTTTGGCTTTG
CTGTCCTGCTTGACGGTTCCAACKACCGCTCACGAGGTGCGCAACGCATCCGGGGTGTATCATGTC
ACCAACGACTGTTCCAACTCGAGCATCATCTATGAGATGGACGGTATGATCATGCACTACCCAGGG
TGCGTGCCCTGCGTTCGGGAGGATAACCATCTCCGCTGCTGGATGGCGCTCACCCCCACGCTTGCG
GTCAAAAAYGCTAGTGTCCCCACTRCGGCAATCCGACGTCACGTCGACTTGCTTGTTGGGGGNNCC
ACGTTCTGTTCCGCTATGTACGTGGGRGACCTTTGCGGGGTCTGTCTCCTCGCTGGCCAGCTATTC
ACCTTTTCACCCCGCATGCACCATACAACGCAGGAGTGCAACTGCTCAATC

SEQ ID NO. 5 (BNL2, 1d)
ATGAGCACGATCCTAAACCTCAAAGAAAAACCAAACGTAACACCAACCGCCGCCCACAGGACGTC
AAGNTCCCGGGTGGTCAGATCGTTGGTGGAGTTTACCTGTTGCCGCGCAGGGGCCCCAGGTTG
GGTGTGCGCGGACCAGGAAGACTTCCGAGCGGTCGCAGCCTCGTGACAGGCGACAGCCTATTCCT
AAGGCTCGCCAGTCCGATGGCAGNNCCTGGGCTCAGCCAGGGCATCCCTGGCCCCTCTATGGCAAT
GAGGGCTGCGGATGGCGGATGGCTCCTGTCCCCCCGCGGCTCTCGGCCCAGTTGGGGCCCC

SEQ ID NO. 7 (BNL2, 1d)
GACGGCGTGAACTATGCAACAGGGAATTTGCCTGGTTGCTCTTTCTCTATCTTCTCTTTAGCTTTT
CTGTCCTGCTTGACGGTTCCAACTACCGCTCATGAGGTGCGCAACGCATCCGGGGTATATCATCTC
ACCAATGACTGTTCCAACTCGAGCATCATCTATGAGATGAGTGGTATGATCTTGCACGCCCCAGGG
TGTGTGCCCTGCGTTCGGGAGAACAACTCTTCTCGTTGCTGGATGCCRCTCACCCCCACGCTTGCG
GTCAAAGACGCTAATGTCCCTACTGCGGCAATCCGACGCCATGTCGACTTGCTGGTTGGGACAGCC
GCGTTTCGTTCCGCTATGTACGTGGGGGACCTCTGCGGATCCGTCTTCCTTGTCGGCCAGCTATTC
ACCTTTTCACCCCCGCTTGTACCATACAACACAGGAGTGCAACTGCTCAATC

Figure 3 - continued

SEQ ID NO. 11 (FR2, 1f) ATGAGCACGAATCCTAAACCTCAAAGAAAAACCAAACGCAACACCAACCGCCGCCCACAGGACGTT AAATTCCCGGGTGGGGGCAGATCGTGGGTGGAGTTTACTTGTTGCCGCGCAGGGCCCCAGGTTG GGTGTGCGCGCGACGAGGAAGACTTCCGAGCGGTCGCAACCTCGCGGAAGGC GACAGCCTATCCCCAAGGCTCGCCGACCCGAGGGCAGGTCCTGGGCTCAGCCTGGGTACC CATGGCCCCTCTATGCTAACGAGGGCTGCGGATGGGCGGGATGGCTCCTGTCCCCTCGCG GCTCCCGTCCTAGCTGGGGCCCCAATGACCCCCGACGTAGATCACGCAATTTGGGTAAGG TCATCGATACCCTAACGTGTGGCTTCGCCGATCTCATGGGGTACATTCCGCTCGTCGGCGC CCCCCTAGGGGGCGCTTCCAGAACCCTGNCACATGGTGTCCGGGTCCTGGNAGGCGGCGTGATNNN NNNNNNNNAACCTTCCNGGTTGCTCTTTNNCTATCTTCCTCTTGGCNTTACTCTCTTGCCTCAC AGTCCCCACCTCTGCCTATGAGGTGCACAGCACAACCGATGGCTACCATGTCACTAATGACTGTTC CAACGGCAGCATCGTATATGAGGCAAAGGACATCATCCTTCACACGCCTGGGTGNGTGCCCTGCAT ACGGGAAGGCAATATCTCCCGTTGCTGGGTACCGCTCACCCCCACGCTCGCAGCGCGGATCGCGAA CGCTCCCATCGATGAGGTGCGGCGTCACGTCGACCTCCTCGTGGGGGCAGCCGTGTTCTGCTCAGC CATGTACATTGGGGACCTTTGTGGGGGGCGTCTTCCTCGTTGGGCAATTGTTCACCTTCACGTCCCG GCGGCATTGGACGGTGCAGGACTGTAATTGTTCCATTTACTCTGGCCACATAACGGGCCACCGNNN NNNN

SEQ ID NO. 13 (BNL3, 2e)
ATGAGCACAAATCCTAAACCTCAAAGAAAAACCAAAAGAAATACCAACCGCCGCCCACAGGACGTC
AAGTTCCCGGGCGGCGGCCAGATCGTTGGCGGAGTTTACTTGTTGCCGCGCAGGGGCCCCAGATTG
GGTGTGCGCGCGACGAGAAAGACTTCTGAACGGTCCCAGCCACGTGGAAGGCGCCCAGCCCATCCCT
AAAGATCGGNGNGCCACTGGCAGGTCCTGGGGACGTCCAGGATATCCCTGGCCCCTGTATGGGAAC
GAGGGGCTCGGCTGGGCAGGATGCCTCCTGTCCCCCCGAGGCTCCTC

SEQ ID NO. 17 (FR4, 2f) ATGAGCACAAATCCTAAACCTCAAAGAAAAACTAAAAGAAACACTAACCGTCGCCCACAGGAC GTTAAGTTCCCGGGCGGCGGCCAGATCGTTGGCGGAGTTTACTTGTTGCCGCGCAGGGGCCCCAG GTTGGGTGTGCGCGCGCCAAGGAAGACTTCTGAACGGTCCCAGCCACGTGGAAGGCGCCAGCCC ATCCCAAAAGATCGGCGCCCCCCTGGCAAGTCCTGGGGACGTCCAGGATACCCTTGGCCCCTGT ACGGGAACGAGGGCCTCGGCTGGGCAGGGTGGCTCCTGTCCCCCCGGGGCTCTCGCCCCTCGTG GGGCCCAAACGACCCCCGGCACAGGTCACGCAACTTGGGTAAGGTCATCGATACCCTCACGTG TGGCTTTGSCGACCTCATGGGGTACATACCTGTCGTCGGCGCCCCTGTGGGCGGCGTTGCCAGA GCCCTCGCGCATGGCGTGCGGGTCCTGGAGGACGGGATAAATTATGCAACAGGGAACTTGCCCGGT GTTAAGAACAACAGCCACTTCTACATGGCGACTAATGACTGTGCCAATGACAGCATCGTCTGGCAG CTCAGGGACGCGGTGCTCCATGTTCCTGGATGTGTCCCCTGTGAGAGGTCAGGTAATAGGACCTTC TGTTGGACAGCGGTCTCGCCCAACGTGGCTGTGAGCCGACCTGGTGCTCTCACTAGAGGTCTGCGG GCTCACATTGATACCATCGTGATGTCCGCCACCCTCTGCTCTGCCCTATACATAGGGGACCTATGC GGCGCTGTGATGATAGCAGCGCAAGTTGCCGTCGTCTCACCGCAATACCATACTTTTGTCCAGGAA TGCAACTGCTCCATATACCCAGGCCATATCACAGGACATCGAATGGNN

Figure 3 - continued

SEQ ID NO. 19 (BNL4, 2g)
GACGGGGTAAATTATGCAACAGGGAATCTGCCTGGTTGCTCTTTCTCTATCTTCTTGTTGGCTCTT
GACGGGGTAAATTATGCAACAGGGAATCTGCCTGGTTGCTGTTAGAACACCAGTACCATGTACATGGCA
CTGTCTTGTGTCACCGTGCCTGTCTCTGCCGTGCAGGGCGCGGGGGGCTTCATGTTCCTGGA
ACCAATGACTGTTCCAACAACAGCATCATCTGGCAAATGCAGGGCGCGGTGCTCACTCCCAACGTGGCT
TGTGTCCCGTGTGAGTTGCAGGGCAATAAGTCCCGGTGCTGATCACCATCGTGATGGTCGCT
GTGAACCAGCCCGGCGCCCTCACTAGGGGCTTGCGGACGCACATTGACACCATCGTGATGGTCGCT
ACGCTCTGTTCTGCACTCTACATCGGGGACGTGTGTGCCAATTGTTCCATC

SEQ ID NO. 21 (BNL5, 2h)
ATGAGCACAAATCCTAAACCTCAAAGAAAAACCAAAAGAAACACTAACCGCCGCCCACAGGACGTT
ATGAGCACAAATCCTAAACCTCAAAGAAAAACCAAAAGAAACACTAACCGCCGCGCCACAGGACGTTG
AAGTTCCCGGGCGCGACGAGAAACTTCCGAACGGTCCCAGCCACGTGGGAGGCGCCAGCCCATCCCT
GGTGTGCGCGCGCCACTGGCAAAACCTTCCGGGACGTCCAGGATACCCTTGGCCCCTGTATGGGAAC
GAGGGCCTTGGTTGGGCAGATGCCTTGTCCCCTCGAGGCTCTC

SEQ ID NO. 23 (BNL5, 2h)
GACGGGATAAACTACGCAACAGGGAATCTGCCCGGTTGCTCCTTTTCTATCTTCTTGCTGGCCTTG
GACGGGATAAACTACGCAACAGGGAATCTGCCCGGTTGCAAGAACACCAGCCACTCTTATATGGTG
CTATCCTGTCTCACTGTGCCGGCGTCCGCTGTGCAGGTCAAGAACACCAGCCACTCTAAACAGCAGCATTGTCTGGCAGCTTAAAGGATGCTGTGCTTCACGTCCCTGGA
ACCAATGATTGCTGAGAGGCACCAAAATCAGTCTCGCTGCTGGATACCTGTGACACCAATGTGGCC
GTGAGCCAACCTGGCGCGCTCACCAGGGGTTTGCGGACGCACATTGACACCATCGTTGCGTCTGCT
ACCGTCTGCTCAGCTTTGTATGTGGGCGACTTCTGCGGCGCAGTGATGTTGGTCTCAATTTTTC
ATGATCTCCCCTCAGCACCACATCTTCGTCCAGGATTGCAACTGCTCGATA

SEQ ID NO. 29 (BNL7, 4k)
GACGGGATCAATTTTGCAACAGGGAACCTCCCCGGTTGCTCCTTTTCTATCTTCCTCTTGGCACTC
GACGGGATCAATTTTGCAACAGGGAACCTCCCCGGTTGCTCCTATGTCTCTCGGGCATTTACTATGTC
CTCTCGTGCCTGACTGTCCCCGCTTCGGCCATCAACTATCGCACTCACATCTTGCACCTCCCAGGT
ACCAATGATTGCCCGAATTCAAGCATAGTGTATGAGGCCGACCATCACATCTTGCACCGTCGCA
TGCGTGCCCTGCGTGAGAGAGGGGAATCAGTCACGTTGCTGGGACTTGATGGTGGGGGGCCGCC
GCGCCATACATCGGCGCCCACTTGAGTCTCTACGGAGTCATGTTCCTAGTCGGTCAGATGTTC
ACTGTTTGTTCAGCCCTTTACATCGGGGATTTRTGTGGGYGGCTTGTTCCATC
TCTTTCCGACCAAGGCGCCACTGGACTACTCAAGATTGCAATTGTTCCATC

Figure 3 - continued

SEQ ID NO. 35 (BNL10, 4k)
GACGGGATCAATTATGCAACAGGGAATATTCCCGGTTGCTCYTTTTCTATCTTCCTTYTGGCACTT
CTCTCGTGTCTGACTGTCCCCGCTTCGGCCACTAACTATCGCAACGTCTCGGGCATCTACCATGTC
ACCAATGACTGCCCGAATTCAAGCATAGTGTATGAGGCCGACCATCACATCTTAGCACTTCCAGGT
TGCGTGCCCTGCGTGAGAGTGGGGAACCAGTCACGCTGCTGGGTGGCCCTTACCCCTACCGTCGCA
GCGCCATACACCGCGGCGCCGCTTGAGTCCCTGCGGAGTCATGTGGATCTGATGGTGGGAGCTGCC
ACTGTTTGTTCAGCCCTTTACATCGGGGAYTTGTGTGGCGGCTTGTTCTTGGTTGGTCAGATGTTC
TCTTTYCAGCCTCGGCGCCACTGGACTACCCAGGATTGCAATTGTTCCATC

Figure 3 - continued

SEQ ID NO. 43 (VN4, 7c) ATGAGCACACTTCCAAAACCCCAAAGAAAAACCAAAAGAAACACCATCCGCCGCCCACA GGACGTCAAGTTCCCGGGTGGCGGCCAGATCGTTGGTGGAGTCTACTTGCTGCCGCGCAG GGGCCCGCGCTTGGGTGTGCGCGCGACGAGAAAGACTTCTGAACGGTCCCAGCCCAGAGG TAGGCGCCAACCAATACCCAAAGTGCGCCACCAAACGGGCCGTACCTGGGCCCAGCCCGG CCGCGGCTCTCGCCCAAATTGGGGCCCAAACGACCCCCGGCGGAGGTCCCGCAACTTGGG TAAAGTCATCGACACCCTTACTTGCGGGCTTCGCCGACCTCATGGGGTATATCCCTGTCGTAG GCGCTCCGWTGGGAGGCGTCGCGGNGGCCTTGGCGCATGGGGTCANGGNCATCGAGGACGGNGTAA ATTACGCAACAGNGAATCTTCCCGGNNGCTCTNTCTCTATCTTNCTCTTGGCACTTCTCTCGTGCC TTACAACACCAGCCTCCGCGGCGCATTATACCAACAAGTCTGGCCTGTACCATCTCACCAACGACT GCCCCAACAGCAGCATCGTTTATGAGGCGGAGACACTGATTTTGCACTTGCCTGGGTGTGTACCTT GTGTGAAGRTGRACAATCAATCCCGGTGCTGGGTGCAGGCCTCCCCGACCCTGGCAGTGCCGAACG CGTCTACGCCAGTCACCGGGTTCCGCAAACATGTGGACATCATGGTGGGCGCTGCCGCGTTCTGTT CAGCTATGTATGTGGGGGGACCTGTGCGGGGGCCTTTTCCTCGTTGGACAGCTCTTCACGCTCAGGC CTCGGATGCATCAGGTTGTCCAGGAGTGTAACTGTTCCATCTACACAGGGCATATCACTGGACACC GAATGGCA

SEO ID NO. 47 (VN12, 7d) ATGAGCACACTTCCAAAACCCCAAAGAAAACCAAAAGAAACACAAACCGTCGCCCAATGGATGTC AAGTTCCCGGGCGGCGGTCAGATCGTTGGTGGAGTCTACTTGTTACCGCGCAGGGGCCCACGTTTG AAGGTGCGCCAGAACCAAGGCCGAACCTGGGCTCAGCCTGGGTACCCCTGGCCCCTTTATGGGAAC GAGGGCTGCGGCTGGGCTGGCTCTTGTCCCCCCGTGGCTCTCGCCCGGACTGGGGNCCCAAT GACCCCGGNGGAGGTCCCGCAACCTGGGTAAGGTCATCG ACACCCTCACTTGCGGCTTCGCCGACCTCATGGAGTACATCCCTGTCGTTGGCGCCCCCCT TGGAGGCGTTGCGGCGGAACTGGNACATGGTGTCAGGGCCATCGAGGACGGGATAAACTATGCAAC AGGGAATCTTCCTGGTTGCTCTTTCTCTATCTTCCWCTTGGCACTTCTCTCGTGCCTCACCACGCC TGCCTCCGCACTAAACTATGCTAACAAGTCTGGGCTGTATCATCTAACCAATGACTGCCCCAATAG CAGCATTGTGTATGAGGCGAATGGCATGATCCTGCATCTCCCGGGTTGCGTCCCCTGCGTGAAGAC CGGCAACCTGACCAAGTGTTGGCTGTCGGCCTCCCCGACATTGGCGGTGCAGAATGCGTCGGTGTC CGTGGGCGACTTATGCGGTGGGCTCTTTCTCGTTGGGCAGTTGTTCACGTTCAGACCCAGGATGTA TGAGATCGCCCAGGACTGCAACTGTTCCATCTATGCAGGCCACATCACTGGGCACCGGATGGCG

SEO ID NO. 41 (FR1, 9a) ATGAGCACACTTCCAAAACCCCAAAGAAAAACCAAAAGAAATACTAACCGTCGCCCTATGGAC GTCAAGTTCCCGGGCGGCGGCCAGATCGTTGGTGGAGTTTACTTGTTGCCGCGCAGGGGC CCTCGTTTGGGTGTGCGCGCGACGAGAAAGACCTCCGAACGGTCCCAGCCTAGAGGCAGG CGCCAGCCCATACCAAAGGTACGCCAGCCGACAGGCCGTAGCTGGGGTCAACCCGGCTAC CCTTGGCCCCTTTATGGCAACGAGGGCTGCGGATGGCGGGATGGCTCCTGTCCCCCGC GGGTCTCGTCCTAATTGGGGCCCCAACGACCCCCGGCGAAGGTCCCGCAACTTGGGTAAG GTCATCGATACCCTTACATNCGGNCTAGCCGACCTCATGGGGTACATCCCTGTCCTAGGAGG GCCGCTTGGCGGCGTTGCCGCTGGCGCATGGCGTTAGGGCAATCGAGGACGGGGTCAATTA CGCAACAGGGAATCTTCCTGGTTGCTCCTTTTCTATCTTCCTCTTAGCACTGTTATCGTGCCTCAC TACACCAGCCTCAGCAATTCAAGTCAAGAACGCCTCTGGGATCTACCATCTTACCAATGACTGCTC GAACAACAGCATCGTTTTTGAGGCGGAGACCATGATACTGCATCTTCCAGGTTGTGTCCCATGTAT CAAGGCGGGGAATGAGTCACGATGTTGGCTCCCTGTCTCCCCCACCTTAGCCGTCCCCAACTCATC AGTGCCAATCCACGGGTTTCGCCGACACGTAGACCTCCTCGTTGGGGCAGCGGCATTTTGTTCGGC CATGTACATCGGAGACCTCTGTGGTAGCATAATCTTGGTAGGGCAGCTTTTTACTTTCAGGCCTAA GTACCATCAGGTTACCCAGGATTGTAACTGCTCTATNAACNCTGGCCACGTCACGGGACACAGGAT GGCA

Figure 3 - continued

SEQ ID NO. 49 (NE98, 10a)
ATGAGCACACTTCCTAAACCACAAAGAAAAACCAAAAGAAACACCAACC?CCGGCCACAGGACGTT

SEQ ID NO. 51 (NE98, 10a)

SEQ ID NO. 53 (BNL1,1d)

CTCGACAGTTACTGAGAATGACATCCGTGTCGAGGAATCAATATACCAATGTTGTGACTTGGCCCCCGAGGCTCGCAAGGCCATAAAGTCGCTCACCGAGCGCTGTACATCGGGGGCCCYCTAACCAATTCAAAAGGACAGAACTGCGGCTACCGTCGGTGCCGCCAGCGGCGTGCTGACTACCAGCTGCGGCAACCCCTGACATGCTACTTGAAAGCCAGAGCGGCCTGTCGAGCTGCAAAGCTCCGGGACTGCACCATGCTCGTGTGTGCGGGATGACCTTGTCGTTATCTGTGAGAGTGCGGGAGTCGAGGAAGACGCGGCGAACCTTACGAGCT

SEQ ID NO. 55 (BNL2,1d)

CTCGACAGTTACTGAGAACGACATCCGTACCGAGGRATCAATCTATCAATGTTGTGACTTGGCCCC
YGAGGCCCGCAAGGCCATAAAGTCGCTCACCGAGCGGCTGTACGTCGGGGGCCCCCTAACCAATTC
AAAGGGGCAGAACTGCGGCTATCGTCGGTGTCGCGCTAGCGGCGTGCTGACCACCAGCTGCGGCAA
CACCCTCACATGCTACTTGAAAGCCAGGGCGGCCTGTCGAGCTGCAAAGCTCCAGGACTGCACGAT
GCTCGTGTGCGGAGACGACCTTGTCGTTATCTGTGAGAGCGCGGGGAGTCGAGGAGGACGCGGGCGAA
CCTACGAGTC

SEQ ID NO. 57 (FR17,1d)

SEQ ID NO. 59 (CAM1078,1e)

CGTACAGCCTCCAGGACCCCCCTCCCGGGAGAGCCATAGTGGTCTGCGGAACCGGTGAG
TACACCGGAATTGCCAGGACGACCGGGTCCTTTCTTGGATCAACCCGCTCAATGCCTGGA
GATTTGGGCGTGCCCCGCAAGACTGCTAGCCGAGTAGTGTTGGGTCGCGAAAGGCCTTG
TGGTACTGCCTGATAGGGTGCTTGCGAGTGCCCCGGGAGGTCTCGTAGACCGTCCACCAT
GAGCACGAATCCTAAACCTCAAAGAAAAAACCAAAAGAAACACCAACCGCCGCCCACAGGA
CGTCAAGTTCCCGGGCGGTGGCCAGATCGTTGGTGGAGCTCTACGTGCTACCGCGCAGGGG
CCCTAGATTGGGTGCGCGCAGCGCGGAAGACTTCGGAGCGGTCGCAACCTCGTGGGAG
GCGCCAACCTATTCCCAAGGAGCGCCGACCCGAGGGCAGGTCCTGGGCGCAGCCCGGGTA
CCCCTGGCCCCTCTATGGTAACGAGGGCTGCGGGTGGGCAGGTCCTCCCCTCG
CGGCTCCCGTCCTAGTTGGGGTCCTACTGACCCCCGGCGTAGGTCACGCAATTTGGGTAA
GGTCATCGATACCCTCACGTGTTGNTTCGCCGACCTCATGGGGTACATACCG

Figure 3 - continued

SEQ ID NO. 61 (CAM1078,1e)

CTCAACGGTCACTGAAGCTGATATCCGAACAGAGGAGTCCATATACCAATGCTGTGACCTGCACCC CGAAGCACGTGTAGCCATCAAGTCTTTGACTGAAAGGCTGTACGTCGGGGGGCCCTTGACCAATTC CAAAAGGGGAGAACTGCGGCTATCGCAGATGCCGTGCCAGCGGCGTCTTGACAACCAGCTGCGGCAA CACCCTCACCTGCTATATCAAGGCCCTAGCAGCCTGTAGAGCTGCCAAGCTCCAGGACTGCACCAT GCTCGTCTGTGGCGACGACCTGGTCGTGATCTGCGAGAGTGTAGGGACCCAGGAGGATGCGGCGAG CCTGCGAGCC

SEQ ID NO. 63 (FR2, 1f)

NTCAACAGTCACTGAGAGTGATATCCGTACAGAGGAGTCCATCTACCAATGCTGTGATCTAGACCC CGAGGCTCGCAAGGCCATAAGGTCCCTCACAGAGAGGCCTTTATATCGGGGGTCCCTGACAAACTC CGAGGCCAGAACTGCCGCCGATGCCGTGCAAGCCGCGTCCTGACGACTAGCTGCGCCAA AAAAGGGCAGAACTGCGGCTACCGCCGATGCCGAGCCTGTCGAGCTCCAGGATTGCTCAAT CACCCTCACCTGTTACATAAAGGCCAGGGCAGCCTGTCGAGGTCCCAAGCTCCAGGATTCCTCGAN GCTCGTCTGTGGCGACCTTGTCGTTATCTGCGAGATCGAGGGTCCANGAGGATCCGTCGAN NNNNNNNNNN

SEQ ID NO. 65 (FR16,1g)

SEQ ID NO. 67 (FR16,1g)

SEQ ID NO. 69 (BNL3, 2e)

CTCGACAGTCACAGAGAGAGATATAAGNACTGAGGAGTCCATATACCAGGCTTGTTCCTTACCCGA GCAGGCCAGAACTGCCATACACTCATTGACTGAGAGACTCTACGTAGGAGGCCCATGATGAACAG CAAAGGGCAATCCTGCGGATACAGGCATTGCCGCGCCAGCGGAGTGCTCACCACCAGTATGGGGAA TACCATCACGTGCTACATCAAGGCCCTAGCGGCTTGTAAAGCAGCAGGAATAGTGGCCCCCACCAT GCTGGTGTGCGGCGATGACCTAGTTGTCATCTCAGAGAGTCAGGGAGTCGAGGAGGACGACCGGAA

SUBSTITUTE SHEET (RULE 26)

SEQ ID NO. 71 (FR4, 2f)

CTCAACCGTCACAGAGAGGGATATAAGAACTGAGGAGTCCATATACCTGGCCTGCTCCTTACCCGA GCAGGCCCGGACTGCCATACATTCATTAACTGAGAGACTTTACGTGGGAGGGCCCATGATGAACAG CAAAGGGCAGTCCTGCGGATACAGGCGTTGCCGCGCTAGCGGAGTGCTCACCACCAGTATGGGGAA CACCATCACGTGTTATGTGAAAGCCCTCGCAGCTTGTAAAGCTGCGGGCATTGTTGCCCCCCACGAT GCTGGTGTGCGGCGATGACCTGGTTGTCATCTCAGAGAGTCAGGGGGGCTGAGGAGGACGAGCGAAA CCTGAGAGTC

SEQ ID NO. 73 (BNL5, 2h)

CTCAACAGTCGCGGAGAGAGACATCAGGACCGAGGAGTCCATTTACCTTGCCTGCTCCTTACCCGA GCAAGCCCGAACTGCCATACATTCATTGACTGAGAGACTTTACGTAGGAGGGCCCATGATGAACAG CAAGGGACAGTCCTGCGGTTACAGACGTTGCCGCGCCAGCGGAGTGCTCACCACCAGCATGGGGAA TACCATCACATGCTATGTGAAGGCATTAGCTGCCTGCAAAGCTGCAGGCATCGTTGCTCCCACGAT GCTGGTTTGTGGCGACGATCTGGTCATCATCTCAGAGAGTCAGGGAACCGAGGAGGATGAGCGGAA CCTGAGAGTC

SEQ ID NO. 75 (FR13,2k)

CGNACANCCTCCAGGCCCCCCCCCCCCGGGAGAGCCATAGTGGTCTGCGGAACCGGTGAG TACACCGGAATTGCCGGGAAGACTGGGTCCTTTCTTGGATAAACCCACTCTATGCCCGGC CATTTGGGCGTGCCCCCGCAAGACTGCTARCCGAGTAGCGTTGGGTTGCGAAAGGCCTTG TGGTACTGCCTGATAGGGTGCTTGCGAGTGCCCCCGGGAGGTCTCGTAGACCGTGCATCAT GAGCACAAAT CCTAAACCTCAAAGAAAAACCAAAAGAAACACTAACCGCCGCCCACAGGA CGTTAAGTTCCCGGGCGGTGGCCAGATCGTTGGCGGAGTATACTTGTTGCCNTGCAGGGG NCCCAGGTNGNGTNTATGCGCAACGANGAAGACTNCCGAACAGTCCCAGCCACGTGGGAG GCGCCAGCCCATCCCGAAAGATCGGNGCACCACTGGCAAGTCCTGGGGACGTCCAGGATA TCCCTGGCCCCTGTATGGGAACGAGGGCCTCGGGTGGGCAGGGTGGCTCCTGTCCCCCG GGGCTCCCGCCCGTCATGGGGCCCCACGGACCCCCGGCATAGGTCGCGCAACTTGGGTAA GGTCATCGATACCCTCACGTNCGGCTTTNCCGACCTCATGGGGTACATTCCCGTCGTTGG CGCCCCAGTAGGNGGCGTCGCCAGAGCTCTCGCGCATGGCGTGAGAGTCCTGGAGGACGG TCTGTCCTGAATTACCGNGCCAGTTTCTGCTGTGGAAATCAAAAACACCAGMAACACATA CATGGTGACTAACGACTGTTCAAACAGYAGCATCACCTGGCAGCTTNNGNNCGCGGTGCT TCACGTTCCTGGATGCGTCCCCTGTGAACGAGGGCCAACAGTTCCCGGTGCTGGATTCC AGTCACGCCCRACGTAKNCGTGAGCCGACCTGGTGCCCTAACCGAGGGTTTGCGATCGCA CATCGACACCATCGTAGCGTCCGCAACATTTTGTTCTGCCCTCTACATAGGGGATGTATG TGGCGCGATAATGATAGCTGCCCAAGTGGTCATCGTCTCGCCGGAGCATCATCACTTTGT CCAGGACTGTAACTGTTCCATCTACCCGGGCCACATAACGGGGCCTCGTATGTNG

SEQ ID NO. 77 (FR13,2k)

ATCCACAGTCACTGAAAGAGACATCAGAGTTGAAGAGTCCGTTTATCTGTCCTGTTCACTTCCCGA GGAGGCCCGAGCTGCCATACACTCACTAACTGAGAGGCTGTACGTGGGAGGTCCCATGCAGAACAG CAAGGGGCAATCCTGCGGATACAGGCGCTGCCGCGCCAGCGGGGTGCTCACCACTAGCATGGGGAA TACTCTCACATGCTACTTGAAGGCCCAGGCGGCCTGCAGGGCCGCGGGCATTGTTGCACCCACAAT GCTGGTGTGTGGCGACGACCTGGTCGTCATCTCAGAGAGTCAGGGGACTGAGAGGGACGAGAACAA CCTGAGACCT

SEQ ID NO. 79 (FR18,21)

CTCAACAGTCACGGAGAGGGACATCAGGAATGAGGAGTCCATATTCCTGGCCTGCTCGTTGCCCGAGGAGGCCCGGACTGATACATTCGCTCACTGAGAGACTCTACATAGGCGGGCCGATGATGAACAGCAAAGGCCAGTCCTGTGGATACAGGCGTTGTCGCGCCAGCGGGGTGTTCACCACTAGCATGGGCAATACCATCACGTGCTATGTGAAAGCCATGGCAGCTTGCAGAGCTGCCGGGATTGACGCCCCCACAATGTTGGTATGTGGCGCGCCTGGTGGTCATCTCAGAGAGTCAGGGGACCGAGGAGGACGAGCGAAATCCTGAGAGTC

SEQ ID NO. 81 (PAK64,3g)

CTCTTGACTCTACTGTCACTGAACAGGATATCAGGGTAGAAGAAGAAATATACCAATGTTGTGACC
TTGAGCCGGAGGCTAGACGGCCAATCAAATCGCTCACGGAACGGCTTTACGTTGGAGGTCCCATGT
TCAACAGCAAGGGGCTCAAATGCGGATATCGCCGTTGCCGTGCTAGCGGTGTATTGCCCACTAGCT
ACGGTAATACAATCACCTGCTACATCAAGGCCCAGAGCGGCTGCTCGAGCTGCGGGCCTTCAAGACC
CATCATTCCTTGTCTGCGGAGATGATTTGGTGGTAGTGGCTGAGAGTTGCGKCGTTGATGAGGAGG
ATAGGGCAGC

SEQ ID NO. 83 (BNL8, 4k)

CTCCACTGTAACCGAAAAGGACATCAGGCCCGAGGAAGAGGTCTATCAGTGTTGTGACCTGGAGCCCGAAGCTCGCAAGGTCTACCGAAACAGCGAAGCTCGCAAAGAAGACTCTACGTGGGCGCCCCATGCACAACAGCAAGGGGAGACCTTTGTGGGTATCGGAAACAGCGAGGGGGGGCTGAGAGCCAGCTTCGGAAACACTGACGTGCTACCTCAAAGCCTCAGCTGCTATTAGAGCGGCAGGGCTGAGAGACTGCACCATGCTGGTTTGCGGTGACACGACTGCACCATGCTGGTTTGCGGTGACGACTGGTCGTCATCGCTGAGAGCGATGGCGTAGAGGGAGATAACCGAGCCCCCAAGCC

SEQ ID NO. 85 (BNL12,41)

CTCCACGTGACTGAAAAGGACATCAGGGTCGAGGAAGAGATCTATCAATGTTGTGACCTGGARCC CGAAGCCCGCAAAGCAATATCCGCCCTCACAGAGAGRCTCTACTTGGGCGGCCCCATGTATAACAG CAAAGGGGAGCTCTGCGGGTATCGGAGGTGCCGCGCGAGCGGAGTGTACACCACAAGTTTCGGGAA CACAGTGACCTGCTATCTTAAGGCCACCGCAGCTACCAGGGCTGCAGGCCTAAAAGACTGCACCAT GCTGGTCTGCGGTGACGACTTGGTCGTCATCGCCGAGAGCGAGGGCGTAGAGGAGGATTCCCAACC CCTCCGAGCC

SEQ ID NO. 87 (EG81, 4m)

CTCCACCGTAACCGAAAGGGACATCAGGGTCGAGGAGGAGGTCTATCAGTGTTGTGATCTGGAGCC AGAGGCCCGCAAGGCAATATCCGCCCTCACGGAGAGACTCTATGTGGGCGGTCCCATGTTTAACAG CAAGGGAGACCTATGTGGCTACCGCAGGTGCCGCGAAGCGGCGTCTACACCACCAGCTTCGGAAA CACACTGACCTGCACAGGCCACGGCCGCCTACCAGAGCGCCGGCCTGAAGGATTGCACAAT GCTGGTTTGCGGGACGACCTGGTCATCGCAGAGAGCGATGGCGTGGACGAGACCGCCGAGC CCTCCAAGCT

SEQ ID NO. 89 (VN13,7a)

CTCAACAGTCACAGAGCGCGATGTCCAGACGGAGCATGACATCTACCAGTGCTGTAAGTTGGAGCC CGCAGCACGGACAGCCATCACATCGCTTACTGACCGATTGTACTNCGGTGGTCCCATGTNTAACTC TAAAGGTCAGGCATGTGGATACCGTAGGTGCAGGGCCAGTGGCGGTCTTGACCACCATCCTGGCCAA TACTCTGACTTGCTACTTGAAAGCTCAGGCGGCATGCAGAGCTGCCGGGCTGAAGGACTTTGACAT GTTGGTCTGCGGAGACGACCTTGTCGTTATTTCGGAGAGTTTGGGGGTCTCGGAGGACACTAGTGC ACTGCGAGCT

SEQ ID NO. 91 (VN4,7c)

CTCGACAGTCACCGAGCGCGACATCCRCACCGAGCACGACATCTACCAATGCTGCCAACTTGACCCGTGGCACGCAACGCGACATCCACCGAGCACGCAACGCCAACGCAACTCAACATCACCCCAAGCATGAACTCCGTGGCAACATCATCTGTGGATACCGTAGGTGCCGAGCCAGTGGCGTGCTCACCACGAGCTTGGGCAATACCCTAACATGCTATTTGAAAGCACAAGCAGCGTGTAGGGCAGAAAGCTCAAAAAACTATGACATGTTAGTCTTGGCGGAGAGGAGTCTCTGAGGATGTTGACGCGTTAGTCTGCGGAGAGCACAAGCTCTGAGGATGTTGACGCCCTGCGAGCA

SEQ ID NO. 93 (VN12,7d)

SEQ ID NO. 95 (FR1,9a)

ATCCACAGTCACGGGGCGCGACATACGCACAGAACNAGACATTTACCTGTCCTGCCAGCTCGACCC
AGAGGCCCGGAAAGCCATAAAGTCTCTCACTGAGAGGCTCTATGTCGGGGGCCCTATGTACAACTC
AAAGGGCCAACTCTGTGGTCAACGCCGATGCCGAGCAAGCGGAGTACTCCCCACAAGCATGGGTAA
CACCATCACATGCTTCCTGAAGGCAACCGCCGCTTGCCGAGCAGCCGGCTTTACAGATTATGACAT
GTTGGTCTGCGGAGACGATTTGGTTGTCGTAACTGAGAGTGCTGGAGTCAACGAGGATATCGCTAA
CCTGCGAGCC

SEQ ID NO. 97 (NE98,10a)

SEQ ID NO. 99 (FR14,11a)

SEQ ID NO. 101 (FR15,11a)

Figure 3 - continued

SEQ ID NO. 103 (FR19,11a)
CGTACAGCCTCCAGGACCCCCCTCCCGGGAGAGCCATAGTGGTCTGCGGAACCGGTGAGTACACC
GGAATTGCCGGGAAGACTGGGTCCTTTCTTGGATTAACCCACTCTATGCCCGGAGATTTGGGCGTG
CCCCCGCAAGACTGCTAGCCGAGTAGCGTTGGGTTGCGAAAGGCCTTGTGGTACTGCCTGATAGGG
TGCTTGCGAGTGCCCCGGGAGGTCTCGTAGACCGTGCACCATGAGCACGAATCCTAAACCTCAAAG
ACAAACCAAAAGAAACACCAACCGCCGCCCACAGGACGTTAAGTTCCCGGGCGGTGGCCAGATCGT
TGGCGGGGTGTACTTGTTGCCGCGCAGGGGCCCCAACCTATCCCCAAGGTTAGGCGCACCACCGCCGTT
GGAGCGGTCCCAGCCGCGTGGGAGGCCCCAACCTATCCCCAAGGTTAGGCGCACCACCGGCCGTT

SEQ ID NO. 105 (FR19,11a)
CTCTACTGTCACAGAGAGGGATATACGAACAGAGGAATCCATYTATCTGGCTTGTCAATTGCCCGA
AGAGGCCCGGAAGGCCATCAAATCACTGACAGAGAGACTATACGTGGGCGGCCCGATGGAAAACAG
CAAGGGCCAGGCCTGCGGATACAGGCGTTGCCGCGCAAGCGGGGTATTCACCACAAGCTTGGGGAA
CACCATGACTTGTTACATCAAAGCCAAGGCGGCTTGTAAAGCCGCTGGCATTGTTGACCCAGTGAT
GCTCGTGTGCGGCGACGACCTAGTGGTCATCTCAGAAAGCAAGGGGGGTGGAGGAGGACCAACGAGA
CCTACGANTC

SEQ ID NO. 2 (BNL1, 1d)
MSTNPKPQRKTKRNTNRRPXXXXXPGGGQIVGGVYLLPRRGPRXGVRATRKTSERSQPRGRRQPIP
KAXRXEGRSWAQPGYPWPLYGNEGCGWAXWLLSPRGSRPNWGP

SEQ ID NO. 4 (BNL1, 1d)
DGVNYATGNLPGCSFSIFLLALLSCLTVPXTAHEVRNASGVYHVTNDCSNSSIIYEMDGMIMHYPG
CVPCVREDNHLRCWMALTPTLAVKXASVPTXAIRRHVDLLVGXXTFCSAMYVXDLCGSVFLAGQLF
TFSPRMHHTTQECNCSI

SEQ ID NO. 6 (BNL2, 1d)
MSTNPKPQRKTKRNTNRRPQDVKXPGGGQIVGGVYLLPRRGPRLGVRATRKTSERSQPRDRRQPIP
KAROSDGXXWAOPGHPWPLYGNEGCGWAGWLLSPRGSRPSWGP

SEQ ID NO. 8 (BNL2, 1d)
DGVNYATGNLPGCSFSIFLLAFLSCLTVPTTAHEVRNASGVYHLTNDCSNSSIIYEMSGMILHAPG
CVPCVRENNSSRCWMXLTPTLAVKDANVPTAAIRRHVDLLVGTAAFRSAMYVGDLCGSVFLVGQLF
TFSPRLYHTTOECNCSI

SEQ ID NO. 10 (CAM1078, le)
MSTNPKPQRKTKRNTNRRPQDVKFPGGGQIVGGVYVLPRRGPRLGVRAARKTSERSQPRGRRQPIP
KERRPEGR

SEQ ID NO. 12 (FR2, 1f)
MSTNPKPQRKTKRNTNRRPQDVKFPGGGQIVGGVYLLPRRGPRLGVRATRKTSERSQPRGRRQPIP
KARRPEGRSWAQPGYPWPLYANEGCGWAGWLLSPRGSRPSWGPNDPRRRSRNLGKVIDTLTCGFAD
LMGYIPLVGAPLGGASRTLXHGVRVLXGGVXXXXXNLXGCSXXIFLLXLLSCLTVPTSAYEVHSTT
DGYHVTNDCSNGSIVYEAKDIILHTPGXVPCIREGNISRCWVPLTPTLAARIANAPIDEVRRHVDL
LVGAAVFCSAMYIGDLCGGVFLVGQLFTFTSRRHWT
VQDCNCSIYSGHITGHXXX

SEQ ID NO. 14 (BNL3, 2e)
MSTNPKPQRKTKRNTNRRPQDVKFPGGGQIVGGVYLLPRRGPRLGVRATRKTSERSQPRGRRQPIP
KDRXATGRSWGRPGYPWPLYGNEGLGWAGWLLSPRGSRPSWG

SEQ ID NO. 16 (BNL3, 2e)
TCXXADLMGYXPVVGAPVGGXARALAXGVRVLEDGINYXTGNLPGCSFSIFXLALLSCVTVPVSXV
EVKNTSQAYMATNDCSNNSIVWQLXDAVLHVPGCVPCENSSGRFHCWIPISPNIAVSKPGALTKGL
RARIDAVVMSATLCSALYVGDVCGAVMIAAQAFIVAPKRHYFVQECNCSIYPGHITGHRMA

Tin,

39/74

Figure 3 - continued

- SEQ ID NO. 18 (FR4, 2f)
 MSTNPKPQRKTKRNTNRRPQDVKFPGGGQIVGGVYLLPRRGPRLGVRAPRKTSERSQPRGRRQPIP
 KDRRATGKSWGRPGYPWPLYGNEGLGWAGWLLSPRGSRPSWGPNDPRHRSRNLGKVIDTLTCGFXD
 LMGYIPVVGAPVGGVARALAHGVRVLEDGINYATGNLPGCSFSIFLLALLSCITVPVSAIQVKNNS
 HFYMATNDCANDSIVWQLRDAVLHVPGCVPCERSGNRTFCWTAVSPNVAVSRPGALTRGLRAHIDT
 IVMSATLCSALYIGDLCGAVMIAAQVAVVSPQYHTFVQECNCSIYPGHITGHRMX
- SEQ ID NO. 20 (BNL4, 2g)
 DGVNYATGNLPGCSFSIFLLALLSCVTVPVSAVQVKNTSTMYMATNDCSNNSIIWQMQGAVLHVPG
 CVPCELQGNKSRCWIPVTPNVAVNQPGALTRGLRTHIDTIVMVATLCSALYIGDVCGAVMIAAQVV
 IVSPQHHNFSQDCNCSI
- SEQ ID NO. 22 (BNL5, 2h)
 MSTNPKPQRKTKRNTNRRPQDVKFPGGGRSLAEYTCARRGKLRRSSMG
- SEQ ID NO. 24 (BNL5, 2h)
 DGINYATGNLPGCSFSIFLLALLSCLTVPASAVQVKNTSHSYMVTNDCSNSSIVWQLKDAVLHVPG
 CVPCERHQNQSRCWIPVTPNVAVSQPGALTRGLRTHIDTIVASATVCSALYVGDFCGAVMLVSQFF
 MISPQHHIFVQDCNCSI
- SEQ ID NO. 26 (BNL6, 2i)
 DGINYATGNLPGCSFSIFLLALLSCITVPVSAVQVANRSGSYMVTNDCSNSSIVWQLEEAVLHVPG
 CVPCEWKDNTSRCWIPVTPNIAVSQPGAXTKGLRTHIDIIVASATFCSALYV
- SEQ ID NO. 28 (BNL7, 4k)
 MSTNPKPQRKTKRNTNRRPMDVKFPGGGQIVGGVYLLPRRGPRLGVRATRKTSERSQPRGRRQPIP
 KARRSEGRSWAQPGYPWPLYGNEGCGWAXWLLSPRGSRPSWGPNDPRRSR
- SEQ ID NO. 30 (BNL7, 4k)
 DGINFATGNLPGCSFSIFLLALLSCLTVPASAINYRNVSGIYYVTNDCPNSSIVYEADHHILHLPG
 CVPCVREGNQSRCWVALTPTVAAPYIGAPLESLRSHVDLMVGAATVCSALYIGDXCXGLFLVGQMF
 SFRPRRHWTTQDCNCSI
- SEQ ID NO. 32 (BNL8, 4k)
 DGINYATGNLPGCSFSIFLLALLSCLTVPASAINYRNTSGIYHVTNDCPNSSIVYEADHHILHLPG
 CVPCVRTGNQSRCWVALTPTVAAPYIGAPLESLRSHVDLMVGAATVCSALYIGDLCGGLFLVGQMF
 SFRPRRHWTAQDCNCSI
- SEQ ID NO. 34 (BNL9, 4k)
 DGINYATGNLPGCSFSIFLLALLSCLTVPASAINYHNTSGIYHITNDCPNSSIVYEADHHILHLPG
 CVPCVRVGNQSSCWVALTPTIAAPYIGAPLESLRSHVDLMVGAATVCSALYIGDLCGGAFLVGQMF
 SFRPRRHWTTQDCNCSI
- SEQ ID NO. 36 (BNL10, 4k)
 DGINYATGNIPGCXFSIFLXALLSCLTVPASATNYRNVSGIYHVTNDCPNSSIVYEADHHILALPG
 CVPCVRVGNQSRCWVALTPTVAAPYTAAPLESLRSHVDLMVGAATVCSALYIGXLCGGLFLVGQMF
 SXQPRRHWTTQDCNCSI
- SEQ ID NO. 38 (BNL11, 4k)
 DGINYATGXLPGCSFSIFLLALLSCLTVPASATNYRNVSGIYHVTNDCPNSSIVFEADHHILHLPG
 CVPCVKEGNHSRCWVALTPTVAAPYIGAPLESLRSHVDVMVGAATVCSALYIGDLCGGLFLVGQMF
 SFRPRRHWTTQECNCSI
- SEQ ID NO. 40 (BNL12, 41)
 DGINYATGNLPGCSFSIFILALLSCLTVPASAQHYRNVSGIYHVTNDCPNSSIVYESDHHILHLPG
 CVPCVKTGNTSRCWVALTPTVAAPILSAPLMSVRRHVDLMVGAATLSSALYVGDLCGGAFLVGQMF
 TFQPRRHWTVQDCNCSI

SUBSTITUTE SHEET (RULE 26)

SEQ ID NO. 46 (VN13, 7a)

MSTLPKPQRKTKRNTNRRPQDVKFPGGGQIVGGVYLLPRRGPRLGVRATRKTSERSQPRGRRQPIP KVRHQTGRTWAQPGYPWPLYGNEGCGWAGWLLSPXGSRPNWGPNDPRXRSRNLGKVIDTLTXXFAD LIEYI

SEQ ID NO. 44 (VN4, 7c)

MSTLPKPQRKTKRNTIRRPQDVKFPGGGQIVGGVYLLPRRGPRLGVRATRKTSERSQPRGRRQPIP KVRHQTGRTWAQPGYPWPLYGNEGCGWAGWLLSPRGSRPNWGPNDPRRSRNLGKVIDTLTCGFAD LMGYIPVVGAPXGGVAXALAHGVXXIEDXVNYATXNLPXXSXSIXLLALLSCLTTPASAAHYTNKS GLYHLTNDCPNSSIVYEAETLILHLPGCVPCVKXXNQSRCWVQASPTLAVPNASTPVTGFRKHVDI MVGAAAFCSAMYVGDLCGGLFLVGQLFTLRPRMHQVVQECNCSIYTGHITGHRMA

SEQ ID No. 48 (VN12, 7d)

MSTLPKPQRKTKRNTNRRPMDVKFPGGGQIVGGVYLLPRRGPRLGVRATRKTSERSQARGRRQPIP KVRQNQGRTWAQPGYPWPLYGNEGCGWAGWLLSPRGSRPDWXPNDPRXRSRNLGKVIDTLTCGFAD LMEYIPVVGAPLGGVAAELXHGVRAIEDGINYATGNLPGCSFSIFXLALLSCLTTPASALNYANKS GLYHLTNDCPNSSIVYEANGMILHLPGCVPCVKTGNLTKCWLSASPTLAVQNASVSIRGVREHVDL LVGAAAFCSAMYVGDLCGGLFLVGQLFTFRPRMYEIAQDCNCSIYAGHITGHRMA

SEQ ID NO. 42 (FR1, 9a)

MSTLPKPQRKTKRNTNRRPMDVKFPGGGQIVGGVYLLPRRGPRLGVRATRKTSERSQPRGRRQPIP KVRQPTGRSWGQPGYPWPLYGNEGCGWAGWLLSPRGSRPNWGPNDPRRSRNLGKVIDTLTXXLAD LMGYIPVLGGPLGGVAAALAHGVRAIEDGVNYATGNLPGCSFSIFLLALLSCLTTPASAIQVKNAS GIYHLTNDCSNNSIVFEAETMILHLPGCVPCIKAGNESRCWLPVSPTLAVPNSSVPIHGFRRHVDL LVGAAAFCSAMYIGDLCGSIILVGQLFTFRPKYHQVTQDCNCSXNXGHVTGHRMA

SEQ ID NO. 50 (NE98, 10a)

MSTLPKPQRKTKRNTNXRPQDVKFPGGGQIVGGVYVLPRRGPQLGVRAVRKTSERSQPRSRRQPIPRARRTEGRSWAQPGYPWPLYGNEGCGWAGWLLSPRGSRPSWGPNDPRRR

SEQ ID No. 52 (NE98, 10a)

DGINFATGNLPGCSFSIFLLALFSCLLTPTAGLEYRNASGLYMVTNDCSNGSIVYEAGDIILHLPGCVPCVRSGNTSRCWIPVSXTVAVKSPCAATASLRTHVDMMVXAATLCSALYVGDLCGALFLXGQGFSWRHRQHWTVQDCNCSI

SEQ ID NO. 54 (BNL1,1d)

STVTENDIRVEESIYQCCDLAPEARKAIKSLTERLYIGGXLTNSKGQNCGYRRCRASGVLTTSCGN TLTCYLKARAACRAAKLRDCTMLVCGDDLVVICESAGVEEDAANLRA

SEQ ID NO. 56 (BNL2,1d)

STVTENDIRTEXSIYQCCDLAXEARKAIKSLTERLYVGGPLTNSKGQNCGYRRCRASGVLTTSCGN TLTCYLKARAACRAAKLQDCTMLVCGDDLVVICESAGVEEDAANLRV

SEQ ID NO. 58 (FR17,1d)

STVTENDIRVEESIYQCCDLAPEARKAIKSLTERLYIGGPLTNSKGQNCGYRRCRASGVLTTSCGN TLTCYLKARAACRAAKLQDCTMLVCGDDLVVICESXGVEEDAANLRV

Figure 3 - continued

SEQ ID NO. 60 (CAM1078, le)

MSTNPKPQRKTKRNTNRRPQDVKFPGGGQIVGGVYVLPRRGPRLGVRAARKTSERSQPRGRRQPIP KERRPEGRSWAQPGYPWPLYGNEGCGWAGXLLSPRGSRPSWGPTDPRRRSRNLGKVIDTLTCXFAD LMGYIP

SEQ ID NO. 62 (CAM1078, le)

STVTEADIRTEESIYQCCDLHPEARVAIKSLTERLYVGGPLTNSKGENCGYRRCRASGVLTTSCGN TLTCYIKALAACRAAKLQDCTMLVCGDDLVVICESVGTQEDAASLRA

SEQ ID NO. 64 (FR2, 1f)

STVTESDIRTEESIYQCCDLDPEARKAIRSLTERLYIGGPLTNSKGQNCGYRRCRASGVLTTSCGN TLTCYIKARAACRAAKLODCSMLVCGDDLVVICEIEGXXEDPSXXXX

SEQ ID NO. 66 (FR16,1q)

MSTNPKPQRKTKRNINRRPQDVKFPGGGQIVGGVYLLPRRGPRLGVRATRKTSERSQPRGRRQPIP KARRSEGRSWAQPGYPWPLYGNEGMGWAGWLLSPHGSRPSWGPSDPRRRSRNLGKVIDTLTCGFAD LMGYIPLVGAPLGGVARALAOGFRDL

SEQ ID NO. 68 (FR16, 1g)

XXVTESDIRVEXSIYQCCDLAPEARVAIKSLTERLYVGGPLTNSKGQNCGYRRCRASGVLTTSCGN TLTCYLKAAAACRAAKLRECTMLVCGDDLVVICESAGVOEDAASXXX

SEQ ID NO. 70 (BNL3, 2e)

STVTERDIXTEESIYQACSLPEQARTAIHSLTERLYVGGPMMNSKGQSCGYRHCRASGVLTTSMGN TITCYIKALAACKAAGIVAPTMLVCGDDLVVISESQGVEEDDRNLXX

SEQ ID No. 72 (FR4, 2f)

STVTERDIRTEESIYLACSLPEQARTAIHSLTERLYVGGPMMNSKGQSCGYRRCRASGVLTTSMGN TITCYVKALAACKAAGIVAPTMLVCGDDLVVISESQGAEEDERNLRV

SEQ ID NO. 74 (BNL5, 2h)

STVAERDIRTEESIYLACSLPEQARTAIHSLTERLYVGGPMMNSKGQSCGYRRCRASGVLTTSMGN TITCYVKALAACKAAGIVAPTMLVCGDDLVIISESQGTEEDERNLRV

SEQ ID NO. 76 (FR13, 2k)

MSTNPKPQRKTKRNTNRRPQDVKFPGGGQIVGGVYLLXCRXPRXXXCATXKTXEQSQPRGRRQPIP KDRXTTGKSWGRPGYPWPLYGNEGLGWAGWLLSPRGSRPSWGPTDPRHRSRNLGKVIDTLTXGFXD LMGYIPVVGAPVXGVARALAHGVRVLEDGINYETGNLPGCSFSISLLALLSITXPVSAVEIKNTXN TYMVTNDCSNXSITWQLXXAVLHVPGCVPCEREGNSSRCWIPVTPXVXVSRPGALTEGLRSHIDTI VASATFCSALYIGDVCGAIMIAAQVVIVSPEHHHFVQDCNCSIYPGHITGPRMX

SEQ ID NO. 78 (FR13, 2k)

STVTERDIRVEESVYLSCSLPEEARAAIHSLTERLYVGGPMQNSKGQSCGYRRCRASGVLTTSMGN TLTCYLKAQAACRAAGIVAPTMLVCGDDLVVISESQGTERDENNLRP

Figure 3 - continued

SEO ID NO. 80 (FR18,21)

STVTERDIRNEESIFLACSLPEEARTVIHSLTERLYIGGPMMNSKGQSCGYRRCRASGVFTTSMGN TITCYVKAMAACRAAGIDAPTMLVCGDDLVVISESQGTEEDERNLRV

SEO ID NO. 82 (PAK64,3g)

STVTEQDIRVEEEIYQCCDLEPEARRAIKSLTERLYVGGPMFNSKGLKCGYRRCRASGVLPTSYGN TITCYIKARAAARAAGLQDPSFLVCGDDLVVVAESCXVDEEDRAALR

SEQ ID NO. 84 (BNL8, 4k)

STVTEKDIR PEEEVYQCCDLE PEARKVITALTERLYVGG PMHNSKGDLCGYRRCRASGVYTTSFGN TLTCYLKASAAIRAAGLRDCTMLVCGDDLVVIAESDGVEEDNRALXA

SEO ID NO. 86 (BNL12,41)

STVTEKDIRVEEEIYQCCDLXPEARKAISALTEXLYLGGPMYNSKGELCGYRRCRASGVYTTSFGN TVTCYLKATAATRAAGLKDCTMLVCGDDLVVIAESEGVEEDSQPLRA

SEQ ID NO. 88 (EG81,4m)

STVTERDIRVEEEVYQCCDLEPEARKAISALTERLYVGGPMFNSKGDLCGYRRCRASGVYTTSFGN TLTCYLKATAATRAAGLKDCTMLVCGDDLVVIAESDGVDEDRRALQA

SEQ ID NO. 90 (VN13,7a)

STVTERDVQTEHDIYQCCKLEPAARTAITSLTDRLYXGGPMXNSKGQACGYRRCRASGVLTTILAN TLTCYLKAQAACRAAGLKDFDMLVCGDDLVVISESLGVSEDTSALRA

SEQ ID NO. 92 (VN4,7c)

STVTERDIXTEHDIYQCCQLDPVARKAITSLTERLYCXGPMMNSRGQSCGYRRCRASGVLTTSLGN TLTCYLKAQAACRAAKLKNYDMLVCGDDLVVIAESGGVSEDVDALRA

SEO ID NO. 94 (VN12,7d)

SSVTERDIRTEHDIYQCCQLDPVARKAITSLTERLYCGGPMYNSRGQSCGYRRCRASGVFTTSLGN TMTCYLKAQAACRAXKLKNFDMLVCGDDLVVIAESGGVPEDAGALRV

SEQ ID NO. 96 (FR1,9a)

STVTGRDIRTEXDIYLSCQLDPEARKAIKSLTERLYVGGPMYNSKGQLCGQRRCRASGVLPTSMGN TITCFLKATAACRAAGFTDYDMLVCGDDLVVVTESAGVNEDIANLRA

SEQ ID NO. 98 (NE98,10a)

STVTEQDIRVELSIFQACDLKDEARRVITSLTERLYCGGPMFNSKGQHCGYRRCRASGVLPTSFGN TITCYIKAKAATKAAGIKNPSFLVCGDDLVVIAESAGIDEDKSALRA

SEQ ID NO. 100 (FR14,11a)

STVTERDIRTEESIYLSCQLPEEARKAIKSLTERLYVGGPMENSKGQACGYRRCRASGVFTTSLGN TMTCYIKAKAACKAAGIVDPVMLVCGDDLVVISESKGVEEDQRDLRV

43/74

SEQ ID NO. 102 (FR15, 11a)

STVTERDIRTEESIXXACQLPEEARKAIKSLTERLYVGGPMENSKGQACGYRRCRASGVFTTSLGN TMTCYIKAXAACKXAGIVDPVMLVCGDDLVVISESKGVEEDQRDLXX

SEQ ID NO. 104 (FR19, 11a)

 ${\tt MSTNPKPQRQTKRNTNRRPQDVKFPGGGQIVGGVYLLPRRGPRVGVRATRKTSERSQPRGRRQPIPKVRRTTGR}$

SEQ ID NO. 106 (FR19,11a)


STVTERDIRTEESXYLACQLPEEARKAIKSLTERLYVGGPMENSKGQACGYRRCRASGVFTTSLGN TMTCYIKAKAACKAAGIVDPVMLVCGDDLVVISESKGVEEDQRDLRX

44/74


Core/El amino acid alignment	SEQ	MSTNPKP	1d 2	A	le 10/60R-TA-				2cR-T	X		18R	76R		I	<u>L</u>		4dR-TM	4eR-TM		4.2			78	Z	LR	46I	44L	d 48LR	a 42LR	10a 50	
- 1			۰ '	د،	9/	12	99	•	•	•	14	18			•	•	•	•	•	•			•	28	, ,	•	46	·	•		4	
ore/E	Type	1a	3 5	1 p	ъ.	₩ (٦ (ر ب	2p	ςς	2d	, 9	7. 7.	2k	3a	3a	3p	4°C	4 d	4 e										7d	9 8	10a 11a	
Figure 4. Co	Isolate	HCV-1	BNI.1	BNL2	CAM1078	FR2	FRIO HC-J6	HC-38				III.					GB358		T CAM600			HPCCOREZB	GB724	12/28 RNI,7	BE95	HK2	VN13	VN4	VN12	FRI	NE98 FR19	

51 core-V	SERSOPRGRROPI PKARRPEGRIWAQPGY PWPLYGNEGCGWAGWLL	X	DDOSD-XXHH			SM		DST-KS-GK	LL	II-KS-GKII	D-XATS-GR	LLDAT-KS-GRL	X-QLD-XTT-KS-GRLL		SS	KQ-HLSRSKLL				SSSS	S	-HSS	S	S	SSSS	Q-TS-GAL				AV-QNQ	-L-	SRTS	TTV
SEQ	3	۲)	9	10/60	12	99					14	18	92								28							46	44	48	42	20	104
Type	la 4	19	1d]e	ı£,	1g	2a	5 p	2c	7q	2e	2£	2k	3a	3a	3р	4 0	4q	4 e	4 e	4	4.5	43	42	42	5a	6а	7a	υ	7d		10a	
Isolate	HCV-1	BNL1	BNL2	CAM1078	FR2	FR16	HCJ6	HCJB	· CH610				FR13						CAM600			HPCCOREEZA	HPCCOREZB	HPCCOREZC	GB724	BE95	HK2	VN13	VN4	VN12	FR1	NE98	FR19
											SL	JBS	ITE	TU	TE	SH	1EE	ET ((Rl	JLI	E 2	6)											

101	RGSRPSWGPTDPRRRSRNLGKVIDTLTCGFADLMGYIPLVGAPLGGAARA		N		X09	L-SN	VSH	^	THRIVVV	VV	¦	į	A	\-\X\-\-\-\-\-\-\-\-\-\-\-\-\-\-\-\-	NN	\lambda\lambda	·^	\n\n\n\n\n\n\n\n	^NN	AA	<u>\lambda\lambda</u>	AA		9-1	HN	XNXXXXIE	-X^	XE	NVL-GV-A-	
SEQ			ر 4 ،	•	_	12	99						18	96									28			46	44	48	42	20
Type	1a	1p	n,	D I]e	1£	1g	رع ھ	SP	2 2	5d	2 е	3£	2, 3,	39	4 _C	4 d	4 e	4 e	4£	49	4h	4 k	5 a	9 9	7a	7c	7d	9a	10a
Isolate	HCV1	HCV-J	BNL 1	BNL2	CAM1078	FR2	FR16	HC-J6	HC-J8				III R4										BNL7	BE95	HK2	VN13	VN4	VN12	FR1	NE98




	I-FAS
864 444 0 4844	25
58 44 44 44 44 44 44 44 44 44 44 44 44 44	10a
BNL11 BNL12 BE95 BE100 HK2 VN4 VN12 FR1	NE98




LAPI	ITDHHLAVS	QILSAPS	QILSAPS	LL-DAMLLVDDR-TH-VL-IPN	LQASL-VPN	LNGMLKTLTKLSASL-VQN	LS-NFETMLIKAELPVSL-VPN	-MS-GG-ILSTIPVSXVKS
40					44	48	42	25
41	4%	5 a	5 a	6a	7c	7d	9a	10a
BNL12	GB724	BE95	BE100	HK2	VN4	VN12	FR1	NE98

25.7	GKLPATQLRRHIDLLVGSATLCSALYVGDLCGSVFLVGQLFTESPRRHWT			VA-VFM-IGT		AV-A-MILS-A-MV	A-V-VI-MVALMIAA-AVIA	AII-MVVAIM-AA-VVVV	TTIIAFIA-M-AS-V-II		-A11-MV1 -TT-MVT	-TTI-AVFA-M-S-F-MIOH-I	TII-AF	TI-A	1 1 1	AF-AR		VGAM-	VMAMF-IG	VGA-LES-S-V-M-A-V-N-N-IGM-S-Q	V-MMAV	-S-VMAAI	BNIBWN	AMIGL	VGA-LEBVMA-V	X-N-MIDII	AVIG	Q-VM	-S-VMAVIX-XGL	S-VMAVIGL	S-VMAVIGA	IAA-LESS-V-MAVI-XGDM-SXQ IGA-LESS-VM-AVIGI,M-S-R	
SEQ) 1	_	r 6 0	12					,	1.0 1.0	200	24	56	97															8	35	ى 10 م	389	
Type	la	<u> </u>	1 1 1	1£	2 a	2p	2°	0	, K	.7 C	2 ₀	2 2 2 2 3	2 <u>i</u>	73 X	3а	39	4 a	4 a	4 P	4c	4c	Δ.	4	27 Z	2 4 D 4	4.	49	4h	쇼 놨.	4, 4 X 5	4. 2 X 7	* *	
Isolate	HCV-1	HCV-J BNT 1	BNL2	FR2	HC-J6	HC-JB	CH610	S83	NE92	BNL3 Fod S	BNI 4				E BR36							GB358	DK13	CAMBOU	GBG 0912 CAMG22	CAMG27	GB549	GB438	BNL7	BNL8	BNL9	BNL11	

LSA-LMSVVMASGAM0	$ ext{VDA-LESFVMAVGAM}$	LGAVTAPAV-Y-A-G-AAALMYRO-A-	FGAVTAPAV-YG-AAALMYRO-A-	ASTGFVA-A-VVSII,A	AST-V-GF-K-V-IMA-AFMGII-BM-01	ASVSIRGV-E-VA-AFM-GIRMVFT	SSV-IHGFVA-AFM-III	PCAATAST-V-MM-XAALXG-SWRH-Q	
41 40					44	48	42	52	
41	4%	5а	5a	6a	7c	7d	9a	10a	
BNL12	GB724	BE95	BE100	HK2	VN4	VN12	FR1	NE98	





44 44 42 52 52

4x 5a 5a 6a 7c 7c 7d 9a

GB724 BE95 BE100 BE100 HK2 VN4 VN12 FR1

SUBSTITUTE SHEET (RULE 26)

Figure	5. NS5E	3 nuc	Figure 5. NS5B nucleotide alignment
Isolate	Type SEQ	SEQ	
HCV-1	<u>,</u>	7	/ 932 / 1887
HCV-J	<u>4</u>		b
BE90	1b		-
BNL1	1d	53	GTATGTCATA
BNL2	1d	55	GTACRATT
FR17	1d	57	GTATGTCATG-
CAM1078	1e	61	AGAGCTTA-AT-CA
FR2	1£	63	NAT-CTTT
IS FR16	1g	6 3	NNNNNNTTGTCRTT
	, p		ACAAA-GTT-CAT-GGG
	5p		ACGGAA-AAAAT-CATGG
ENE3	2 9	69	GAATAA-NTT-CAGG
H FR4	S£	71	ACAGTAA-ATT-CATGG
S BNL5	Sh	73	AG-G $-$ AA-GCT-CT-G
H FR13	رع بخ	77	AT-CG-TT-TG-
FR18	21	79	AGGGA-G-ATT-CA-TTGG
딜 (RI	3а		ATACAGA-GGTAAGA
5E JLI	39		TTACATA-GAGA
PAK64	3g	81	TTACAGTA-GGTAA-A-A-A-A

26)

```
C--A-AA-
                   C--A-AG
    7932
SEQ
ID
                                          89
91
93
97
99
101
                            83
85
87
Type
                 440
446
447
441
770
770
110a
111a
111a
Isolate
      GB48
GB116
GB215
GB358
GB309
                       GB549
BNL8
                                  EG81
CHR18
                               BNL12
                                                VN12
FR1
NE98
FR14
FR15
                                          VN13
VN4
                                            SUBSTITUTE SHEET (RULE 26)
```

7982	rgaccrccaagcccgcgrgccarcaagrccrc	T-G-CA-GCAA-GA-	T-G-CG-GA-ACAAA	T-G-CG-GTAAAG	T-G-CYG-GAAAG	-CT-G-CG-GAAAG	-CGCGATATT-GTA	-CTAG-GTAAA-GA	-CG-CG-G-TAAGT	CTC-T-GCC-GAGG-GA-ACTAC-CAGT	CTCTGCCT-AAGA-AACT-TAC-CGT	CTC-T-ACC-GAGGA-AACTAC-CAT-GT	CCCTC-T-ACC-GAGGGACTAC-TAT-AT	CCCTC-T-ACC-GAGAACTAC-TAT-GT	CCTCATCC-GAGG-GA-CTAC-C-AAT	- 1	-CATAGG-GA-GAGA-TGTCCG	-CTG-AG-GTGAAGTGCG-TA	TGGG-GTA-ACGAAGA	
SEQ				53	55	57	61	. 63	29	•		69	71	73	77	19			81	
Type	1a	1b	15	1q	1d	1q	le 1	1£	1g	7a B	qz	92	2£	Sh	2 k	21	3a	33	3g	
Isolate	HCV-1	HCV-J	BE90	BNL1	BNL2	FR17	CAM1078	FR2		1 HC-J6		ENT3	IN.	H BNL5	HS FR13	3 FR18]]]	ဥ RU	F PAK64	26)

```
8031
                                  --AA--TG--ATCCG-
                                                                 --cc-at-a--t--ggt---a--gaaa-
                                                             .C--CC-A--T----
                                                   CA-TGT--T-GC-G
SEQ
ID
                                                       89
91
93
95
97
101
103
                                     83
85
87
                                                       7a
7c
7d
9a
10a
11a
11a
                                              E
                                                   ď
Isolate
        GB48
GB116
                 GB215
GB358
GB809
GB549
BNLB
BNL12
EG81
CCHR18
                                                                VN12
                                                                     FR1
NE98
                                                                             FR14
                                                            SUBSTITUTE SHEET (RULE 26)
```

8032 AGGCTTTATGTTGGGGCCCTCTTACCAATTCAAGGGGGGAACTGCGG C
SEQ 1D 1D 53 53 63 63 77 77 79
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
Isolate HCV-1 HCV-1 HCV-1 HCV-1 HCV-1 BE90 BNL1 BNL2 FR17 CAM1078 FR2 FR16 HC-J6 HC-J6 HC-J6 HC-J6 HC-J6 HC-J6 HC-J6 HC-J6 HC-J7 FR13 FR13 FR13 FR13 FR13 FR13 FR13 FR13

```
8081
                                                                                --TC-A--TC--TCA--T
                                              -CAGC-A---A--CCTT--7
                                                         -CAGC-A---A--CCTA--
                                                               -CAGC-A---C-AC-A-
                                                                          C--CC-T--TC-ATCA--
                                                                     C--T-AA--TC--GCA--
                                          CCTA-
                                     CCTT-
                                          C--C-A---
                                              CA-GCA--
                                                                          --CA-G-TG-
                  --A--
                        --A--
 SEQ
ID
                                                                                                         105
                                                                                                   101
                                                                  993
93
93
94
                                            83
85
87
                                                             5a
7a
7c
7d
9a
11a
Isolate
          GB48
GB116
                    GB215
GB358
                                GB809
                                     GB549
                                           BNL8
BNL12
EG81
CHR18
                                                                 VN13
                                                                                                  FR15
FR19
                                                                             VN12
                                                                                       NE98
                                                                                             FR14
                                                                       VN4
                                                                                 FR1
                                                                             SUBSTITUTE SHEET (RULE 26)
```

```
8131
CTATCGCAGGTGCCGCGCGGCGCGTACTGACAACTAGCTGTGGTAACA
                        -----C-A--
                                                  --C---C-A-
                                                                               --CA-GC-T-
                                                                   --CA-GC-T-
                                                                         A--CA-GCAT
                                                                                    --CA-AC-T
SEQ
ID
                            53
57
61
63
67
                                                                                                               81
Type
                                      ld
                    BE 90
BNL1
BNL2
FR17
CAM1078
FR2
FR16
HC-J6
HC-J8
Isolate
                HCV-J
                                                                                  BNL5
                                                                                       FR13
                                                                                             FR18
                                                                             FR4
                                                                               SUBSTITUTE SHEET (RULE 26)
```

```
CTAC--
Type SEQ
ID
                                          89
91
93
97
99
101
                            83
85
87
                                  4m
5a
7a
7c
7d
9a
10a
11a
Isolate
      GB48
GB116
GB215
GB215
GB358
GB549
BNL12
EG81
CCHR18
VN13
VN4
VN12
FR1
FR1
                                                              FR15
FR19
                                                SUBSTITUTE SHEET (RULE 26)
```

8181 CCCTCACTTGCTACATCAAGGCCCGGCAGCCCTCCTCTCTCT
SEQ 1D 53 53 53 63 63 77 73 79
Type 11a 11b 11d 11d 12b 12b 13a 33a 33a 33a
Isolate HCV-1 HCV-1 HCV-1 HCV-1 HCV-1 BE90 BNL1 BNL2 FR17 CAM1078 FR2 FR16 HC-J6 HC-J6 HC-J8 BNL3 FR16 HC-J8 BNL3 FR16 HC-J8 BNL3 FR17 FR18 FR19 FR19 FR19 FR19 FR19 FR19 FR19 FR19

```
8181
                        --TCA--C--TATCA----
                                 GTT--G--TAC-A-G
                                                            ---A--A-
     8132
SEQ
ID
                                                      89
91
93
95
99
101
105
                                                      7a
7c
7d
9a
10a
Isolate
        GB48
GB116
GB215
GB358
GB809
GB549
BNL8
BNL12
EG81
CHR18
                                                     VN13
                                                          VN12
VN12
FR1
NE98
                                                                           FR14
FR15
FR19
                                                        SUBSTITUTE SHEET (RULE 26)
```

8231 CAGGACTGCACCATGCTGTGTGGCGACGACTTAGTCGTTATCTGTGA
SEQ 1D 53 55 57 69 77 77 79 81
Туре 11 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
Isolate HCV-1 HCV-1 HCV-1 HCV-1 BE90 BNL1 BNL2 FR16 FR16 HC-J6 HC-J6 HC-J8 BNL3 FR16 HC-J8 FR16 HC-J8 FR16 HC-J8 FR17 FR19 T1 FR13 FR19 T1 FR19 FR19 FR19 FR19 FR19 FR19 FR19 FR1

```
1-AA-T
         AGA---
AGA---
AGA---
SEQ
ID
                                    89
91
93
97
97
101
                        83
85
87
                     Isolate
     GB48
GB116
GB215
GB358
GB809
GB549
BNL8
BNL12
EG81
CHR18
                                      VN4
VN12
FR1
NE98
FR14
FR15
                                       SUBSTITUTE SHEET (RULE 26)
```

```
AAGCGCGGGGTCCAGGAGGACGCGGCGAGCCTGAGAGCC
                         ----AC---I
                                     -A---AC---1
                   ----AC---
                                                                                                                 --AGA--AGCT---
                                                                                   -A-CGA-A--
                                                                        -A-CGA-A--
                                                                                         T-A-CG--A--
                                                                                               -A-AAC-A-
                                                                  --A-CG--A-
                                                                             -ACCG--A-
                                                                                                     -A-CGA-AT
                                                           -T--T--A--
                                                      -TC--T---
                  G--T----AAC----
                              G--T----A---G---A
                                                                                              --TCA----ACTG--AG
                                                                                         --TCA---AAC-G--
                                                                       ---CAA--TAA-G-
                                          T--R--A---(
                                                      --N----N---
                                               r-ta---ac
SEQ
ID
Type
                       Isolate
                                              CAM1078
                                                                             HCV-J
                                                                       HC-JB
                                  BNL2
FR17
                                                          FR16
                       BE90
                             BNL1
                                                    FR2
```

```
8271
                                             ---CGCCGAGC---
                                                      -A-TAGTGCA--
                                        --TT-CCAACC--
                   --AAACGAGC--
                           --AAACGANC--
                                T--AAGAGC--
                                    -TAACCGAGC--
                                                                  -TATC--T-A--
                                                          -T-TT-ACGC--
                                                               --29-9-2--L-
                       --AAACGAGC-
                                                 --TAAA---
SEQ
ID
                                                    89
91
93
95
99
101
Type
                     Isolate
        GB18
GB116
GB215-
GB215-
GB358
GB809
GB549
BNL8
BNL12
EG81
CHR18
                                                              V LE SHEET (RULE 26)
```

e 6. NS5B amino acid alignment	Type SEQ	ID 2645	2694 STVTESDIRTEEATYOCCDI. Den betwater were viscon missensial	1bNS	-U-H		Y		62AS		68 XXV-XSA-R	RHR-S-RA-S-RA-S					72RSLA-S-PEH	74A-RSLA-S-PE	78RVSV-LS-S-PEEA	80RNS-FLA-S-PERTV-H			3a	H	<u>E-E-R</u>
l l	Typ		la	1Þ	1c	.1d	1d	1d	1 e	1£	1g	2a	g2	, 7 C	pz	2e	2£	5h	2,k	21	3a	3a	3a	39	39
Figure 6	Isolate		HCV-1	HCV-J	2 T.X 4	BNL1	BNL2	FR17	CAM1078	FR2	FR16	HC-76	HC-J8		S NE92	ENIT?	ES ES TU	E BNL5	HE FR13		BR34			61 26	PAK64

KVEV		KVEVE	1-11) [E	1 [1] 1 [2] 1 [3]	1 E	PR-X-VEVN-EXDX-KV-NA	14	TU-V	KPEVEVEVK	KVEX-EKSAXIMYKI.	TI II	07-NHHHRSSSHH	XX V III		RXHDOVKICXMMOS)WA		QV-LS-E-AKDERV-TCMFK-OH			RSX-LA-Q-PEEKMEK-OA		
												98										102			
4°C	4 _C	4 C	4 C	4 e	4 £	49	4h	054i	1 4j	4.2	4 k	41	4m	5a	5a	7a	7c	7d	9a	10a	11a	11a	11a		
GB48	GB116	GB215	GB358	CB809	CAMG22	GB549	GB438	CAR4/12	CAR1/50	EG13	BNL8	BNL12	EG81									U) FR15	FR19	E 2	6)

2695	2744 Yrrcrasgvittscgntitcytkaraacraacrongmy	3)TAAUUU99AUUT9GAGAGAGAGAGAGAGAGAGAGAGAGAGAGAGAGAGAGA				XX	XXXXXXX			$\frac{1}{2} = \frac{1}{2} = \frac{1}$	·		IXUI		-SXVXI	XTX	I'0	FKVWIDAP	- !	BSPDF	RNPDF	FIKSK-PSF		
SEQ TD	1			54	56	58	29	64	89					70	72	74	78	80				(28	
Type	la	1p	1°	1q	1q	1q	1e	1£	1g	ري م	2p	22	5d	2e	2£	5h	2k	21	g.	3a	3a	g.	39	
lsolate	HCV-1	HCV-J	2TY4	BNL1	BNL2	FR11	CAM1078	FR2	FR16	HC-J6	HC-18	ARG8	S NE92	BNI3	F.	ENIT?						_	FAK64	6)

```
90
94
96
98
100
100
106
                         GB549 49
GB438 4h
CAR4/12054i
CAR1/501 45
EG13 42
BNL8 4k
                     CAMG22
GB48
GB116
GB215
GB358
GB809
                                                              SUBSTITUTE SHEET (RULE 26)
```

```
2757
SAGVQEDAASIRA
---T---
                                                                                                                                     -CX-D-EDRAALR
                 --T----
                                                                                                  -Q-TER-ENN--F
-Q-TE--ERN--V
                                                               -Q-TE--ERN---
-Q-NE--ERN---
-Q-TE--ERN---
-Q--E--DRN-
                                                                                       Q-AE--ERN--
                                                                                             Q-TE--ERN--
                                         -X--E----N--
                                   --N----B---
                                                                                                                                -C--E--R-A--
                             ---N---
                                                    E-XX--PS
                                               -V-T---
SEQ
ID
                                                                                                                                     82
Type
                                                           333333746 dbag
3333322222333
Isolate
                                              CAM1078
           HCV-1
HCV-J
BE90
BNL1
BNL2
FR17
                                                                                  SUBSTITUTE SHEET (RULE 26)
```

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:
BLACK BORDERS
☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
☐ FADED TEXT OR DRAWING
☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
☐ SKEWED/SLANTED IMAGES
☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
☐ GRAY SCALE DOCUMENTS
☐ LINES OR MARKS ON ORIGINAL DOCUMENT
☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
П отнев.

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)